

<https://helda.helsinki.fi>

---

## Essential coastal habitats for fish in the Baltic Sea

Kraufvelin, Patrik

2018-05-01

---

Kraufvelin , P , Pekcan-Hekim , Z , Bergström , U , Florin , A-B , Lehikoinen , A , Mattila , J , Arula , T , Briekmane , L , Brown , E J , Celmer , Z , Dainys , J , Jokinen , H , Kääriä , P , Kallasvuo , M , Lappalainen , A , Lozys , L , Möller , P , Orio , A , Rohtla , M , Saks , L , Snickars , M , Støttrup , J , Sundblad , G , Taal , I , Ustups , D , Verliin , A , Vetemaa , M , Winkler , H , Wozniczka , A & Olsson , J 2018 , ' Essential coastal habitats for fish in the Baltic Sea ' , Estuarine, Coastal and Shelf Science , vol. 204 , pp. 14-30 . <https://doi.org/10.1016/j.ecss.2018.02.014>

---

<http://hdl.handle.net/10138/311712>

<https://doi.org/10.1016/j.ecss.2018.02.014>

---

cc\_by\_nc\_nd

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*


*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# Accepted Manuscript

Essential coastal habitats for fish in the Baltic Sea

Patrik Kraufvelin, Zeynep Pekcan-Hekim, Ulf Bergström, Ann-Britt Florin, Annukka Lehikoinen, Johanna Mattila, Timo Arula, Laura Briekmane, Elliot John Brown, Zuzanna Celmer, Justas Dainys, Henri Jokinen, Petra Kääriä, Meri Kallasvuo, Antti Lappalainen, Linas Lozys, Peter Möller, Alessandro Orio, Mehis Rohtla, Lauri Saks, Martin Snickars, Josianne Støttrup, Göran Sundblad, Imre Taal, Didzis Ustups, Aare Verliin, Markus Vetemaa, Helmut Winkler, Adam Wozniczka, Jens Olsson



PII: S0272-7714(17)30622-4

DOI: [10.1016/j.ecss.2018.02.014](https://doi.org/10.1016/j.ecss.2018.02.014)

Reference: YECSS 5755

To appear in: *Estuarine, Coastal and Shelf Science*

Received Date: 11 June 2017

Revised Date: 1 December 2017

Accepted Date: 12 February 2018

Please cite this article as: Kraufvelin, P., Pekcan-Hekim, Z., Bergström, U., Florin, A.-B., Lehikoinen, A., Mattila, J., Arula, T., Briekmane, L., Brown, E.J., Celmer, Z., Dainys, J., Jokinen, H., Kääriä, P., Kallasvuo, M., Lappalainen, A., Lozys, L., Möller, P., Orio, A., Rohtla, M., Saks, L., Snickars, M., Støttrup, J., Sundblad, G., Taal, I., Ustups, D., Verliin, A., Vetemaa, M., Winkler, H., Wozniczka, A., Olsson, J., Essential coastal habitats for fish in the Baltic Sea, *Estuarine, Coastal and Shelf Science* (2018), doi: 10.1016/j.ecss.2018.02.014.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Essential coastal habitats for fish in the Baltic Sea

Patrik Kraufvelin<sup>(a,b,\*</sup>, Zeynep Pekcan-Hekim<sup>(a</sup>, Ulf Bergström<sup>(a</sup>, Ann-Britt Florin<sup>(a</sup>,  
Annukka Lehtikainen<sup>(a,c</sup>, Johanna Mattila<sup>(a,b</sup>, Timo Arula<sup>(d</sup>, Laura Briekmane<sup>(e</sup>,  
Elliot John Brown<sup>(f</sup>, Zuzanna Celmer<sup>(g</sup>, Justas Dainys<sup>(h</sup>, Henri Jokinen<sup>(i</sup>, Petra  
Kääriä<sup>(j</sup>, Meri Kallasvuo<sup>(k</sup>, Antti Lappalainen<sup>(k</sup>, Linas Lozys<sup>(h</sup>, Peter Möller<sup>(l</sup>,  
Alessandro Orio<sup>(m</sup>, Mehdi Rohtla<sup>(d</sup>, Lauri Saks<sup>(d</sup>, Martin Snickars<sup>(b</sup>, Josianne  
Støttrup<sup>(f</sup>, Göran Sundblad<sup>(n,o</sup>, Imre Taal<sup>(d</sup>, Didzis Ustups<sup>(e</sup>, Aare Verliin<sup>(d</sup>, Markus  
Vetemaa<sup>(d</sup>, Helmut Winkler<sup>(p</sup>, Adam Wozniczka<sup>(g</sup> & Jens Olsson<sup>(a</sup>

a. Swedish University of Agricultural Sciences, Department of Aquatic Resources,  
Institute of Coastal Research, Skolgatan 6, 74242 Öregrund, Sweden

b. Åbo Akademi University, Environmental and Marine Biology, Turku, Finland

c. University of Helsinki, Kotka Maritime Research Centre, Kotka, Finland

d. Estonian Marine Institute, University of Tartu, Tartu, Estonia

e. Institute of Food Safety, Animal Health and Environment "BIOR", Riga, Latvia

f. DTU Aqua, National Institute of Aquatic Resources, Technical University of Denmark,  
Lyngby, Denmark

g. National Marine Fisheries Research Institute, Gdynia, Poland

h. Nature Research Centre, Vilnius, Lithuania

i. University of Helsinki, Tvärminne Zoological Station, Hanko, Finland

j. HELCOM Secretariat, Helsinki, Finland

k. Natural Resources Institute Finland (Luke), Helsinki, Finland

l. Fisch und Umwelt, Mecklenburg-Vorpommern, Rostock, Germany

m. Swedish University of Agricultural Sciences, Department of Aquatic Resources,  
Institute of Marine Research, Lysekil, Sweden

n. AquaBiota Water Research, Stockholm, Sweden

o. Swedish University of Agricultural Sciences, Department of Aquatic Resources,  
Institute of Freshwater Research, Drottningholm, Sweden

p. University of Rostock, Rostock, Germany

\* corresponding author: [patrik.kraufvelin@slu.se](mailto:patrik.kraufvelin@slu.se); [patrik.kraufvelin@abo.fi](mailto:patrik.kraufvelin@abo.fi)

## ABSTRACT

Many coastal and offshore fish species are highly dependent on specific habitat types for population maintenance. In the Baltic Sea, shallow productive habitats in the coastal zone such as wetlands, vegetated flads/lagoons and sheltered bays as well as more exposed rocky and sandy areas are utilised by fish across many life history stages including spawning, juvenile development, feeding and migration. Although there is general consensus about the critical importance of these essential fish habitats (EFH) for fish production along the coast, direct quantitative evidence for their specific roles in population growth and maintenance is still scarce. Nevertheless, for some coastal

species, indirect evidence exists, and in many cases, sufficient data are also available to carry out further quantitative analyses. As coastal EFH in the Baltic Sea are often found in areas that are highly utilized and valued by humans, they are subjected to many different pressures. While cumulative pressures, such as eutrophication, coastal construction and development, climate change, invasive species and fisheries, impact fish in coastal areas, the conservation coverage for EFH in these areas remains poor. This is mainly due to the fact that historically, fisheries management and nature conservation are not integrated neither in research nor in management in Baltic Sea countries. Setting joint objectives for fisheries management and nature conservation would hence be pivotal for improved protection of EFH in the Baltic Sea. To properly inform management, improvements in the development of monitoring strategies and mapping methodology for EFH are also needed. Stronger international cooperation between Baltic Sea states will facilitate improved management outcomes across ecologically arbitrary boundaries. This is especially important for successful implementation of international agreements and legislative directives such as the Baltic Sea Action Plan, the Marine Strategy Framework Directive, the Habitats Directive, and the Maritime Spatial Planning Directive, but also for improving the communication of information related to coastal EFH among researchers, stakeholders, managers and decision makers. In this paper, efforts are made to characterize coastal EFH in the Baltic Sea, their importance and the threats/pressures they face, as well as their current conservation status, while highlighting knowledge gaps and outlining perspectives for future work in an ecosystem-based management framework.

## 1. INTRODUCTION AND BACKGROUND

Fish are central for the functioning of food webs and ecosystems in the Baltic Sea (Österblom et al. 2007, Östman et al. 2016), and are broadly used in environmental monitoring as indicators for ecosystem status and health (Bergström et al. 2016ab). Fish are also important in socio-economic terms, such as for commercial and recreational fisheries (Holmlund and Hammer 1999). The geographical distribution and occurrence of fish in the Baltic Sea, and thereby the species composition of fish communities, differ over both large and small scales. Fish distributions are largely driven by spatiotemporal differences in natural biotic and abiotic factors as well as by human pressures (Bergström et al. 2016a, Östman et al. 2017). The same habitat may have only one or several functions during different seasons with regard to e.g. spawning, feeding and overwintering for the same or different species (Aro 1989, Vetemaa et al. 2006). Examples of common gradients and factors that are determining fish distribution are salinity, temperature, depth, pollution, eutrophication, predation, food availability, fishing pressure, and also the availability and conditions of coastal essential fish habitats (EFH) which is the focus of this review article (Leppäkoski and Bonsdorff 1989, Sparholt 1994, Bonsdorff and Pearson 1999, MacKenzie et al. 2007, HELCOM 2010, Olsson et al. 2012, Seitz et al. 2014). In the review, efforts are made to characterize coastal EFH in the Baltic Sea, their importance and the threats/pressures they face, as well as their current conservation status, while highlighting knowledge gaps and outlining perspectives for future work in an ecosystem-based management framework.

In a broad sense, an EFH is any environment that is needed for the maintenance of a fish population. More specifically, coastal EFH are defined as shallow and nearshore

waters and substrates necessary to any life-stage of fish for spawning, breeding, feeding or growth to maturity (Benaka et al. 1999, Rosenberg et al. 2000). In this respect, the term *waters* include all aquatic coastal areas (down to a maximum depth of 10–20 m) and their physical, chemical, and biological properties, whereas *substrates* include surfaces and their associated biological communities that make them suitable as fish habitats (Rosenberg et al. 2000). Coastal EFH are thus comprised of juvenile growth areas, foraging areas, reproduction areas and migratory routes. While the latter three are of direct importance for fisheries, by offering high catches or value per fishing effort (Airoldi and Beck 2007, Seitz et al. 2014), the former one is a step required to produce recruits to replenish the fishery (Beck et al. 2001). Fishing may, however, be challenging for the sustainable management of some coastal EFH, not only as some fishing practices are detrimental to the habitats per se, but also because targeted extraction of species from the general marine ecosystem may indirectly influence the habitats by altering predator-prey interactions (Hopkins 2003, Eriksson et al. 2011, Pikitch et al. 2014, Östman et al. 2016, Pommer et al. 2016, Eddy et al. 2017). Despite consensus among scientists on the critical importance of EFH, their role for sustaining fish stocks and communities has received relatively little attention (Beck et al. 2001, Gillanders et al. 2003, Armstrong and Falk-Petersen 2008, Sheaves et al. 2015). The influence of the amount and quality of EFH on fish population dynamics has generally been poorly described in the scientific literature, and only rarely, has the information been incorporated into scientific advice for fisheries management (Mangel et al. 2006, Armstrong and Falk-Petersen 2008, Thrush and Dayton 2010, Kallasvuori et al. 2017). As coastal EFH are often found in areas that are highly valued and utilized by humans (de Groot et al. 2012, Šiaulys et al. 2012), numerous pressures/threats and management

issues are implied (Korpinen et al. 2012) and thus the gaps in knowledge with regard to the importance of coastal EFH need to be addressed (Sundblad and Bergström 2014).

Coastal EFH represent “home grounds” for coastal fish species throughout their lives and for other fish species during different life history stages when they are using the coastal zone. Major coastal EFH consist of: (1) coastal wetlands and shallow bays (including salt marshes, estuaries, river mouths, coastal lagoons and flads), (2) shallow vegetated areas (including seagrass meadows and macroalgal beds, but also freshwater plants in brackish water areas), (3) biogenic reefs and hard structures (including mussel beds, rocky shores, mariculture installations and other artificial substrates) and (4) unvegetated soft and sandy areas and shallow open water (modified from Seitz et al. 2014). Thus, basically, most types of shallow benthic and pelagic areas can function as coastal EFH, at least for some species at some life stage. In temperate waters, shallow and wave-sheltered EFH are generally characterised by higher water temperatures, extensive macrophyte vegetation and a particularly high production of zooplankton and zoobenthic prey, thus providing excellent conditions for survival and growth of fish larvae and juveniles (Blaber and Blaber 1980, Karås and Hudd 1993, Gibson 1994, Karås 1996, Ljunggren 2002, Stål et al. 2007, Härmä et al. 2008, Kallasvuo et al. 2009, Snickars et al. 2009, 2010, Ljunggren et al. 2010, Seitz et al. 2014). Many habitats, such as seagrass and macrophyte meadows, perennial macroalgal belts and mussel beds, also aid in maintaining fish populations by providing three-dimensional benthic structures serving as more or less permanent habitats, temporary nursery areas, rich feeding areas and refuges/shelter from predation (Rajasilta et al. 1989, Jackson E.L. et al. 2001, Pihl and Wennhage 2002, Lappalainen et al. 2004, 2005, 2008, Härmä et al. 2008, Díaz et al. 2015). Mariculture installations, artificial substrates and rocky bottoms, in turn, are



important for providing surfaces for habitat-forming macroalgae and sessile animals, which serve as food and refuge from predation (Pihl and Wennhage 2002, Seaman 2007, Fabi et al. 2011, Kraufvelin and Díaz 2015, Bergström et al. 2016c). Finally, seabeds without macroscopic vegetation as well as open shallow waters are often highly productive, both with regard to primary and secondary production (Gerbersdorf et al. 2005, Engelsen et al. 2008). As such they support a diverse range of fish by providing spawning, juvenile growth, feeding and resting grounds (McCormick et al. 1998, Wennhage and Pihl 2002, Cattrijsse and Hampel 2006, Florin et al. 2009, Seitz et al. 2014).

Despite the increased attention during recent years towards characterizing EFH in the Baltic Sea (HELCOM 2012, Sundblad et al. 2014, Kallasvuo et al. 2017), sufficient information is lacking for many fish species to quantitatively assess the role of coastal habitats for fish population growth and production. In this review, the main focus is on the role of coastal essential habitats for commercial, threatened and ecologically important (from a conservation perspective) fish species. The species and groups that benefit from a decrease in the environmental status of the Baltic Sea, such as cyprinids (Bergström et al. 2016ab) and three-spined stickleback (*Gasterosteus aculeatus*) (Bergström et al. 2015, Byström et al. 2015), are thus excluded. Within this process, the threats to and current conservation status of coastal EFH in the Baltic Sea are also thoroughly reviewed, while knowledge gaps are highlighted and perspectives for future work on this topic within an ecosystem-based management framework are outlined.


## 2. OCCURRENCE AND IMPORTANCE OF COASTAL EFH IN THE BALTIC SEA

### 2.1. Occurrence of coastal EFH in the Baltic Sea

The Baltic Sea is the world's largest semi-enclosed brackish water area, with a surface salinity gradient ranging from 2 in the northern and easternmost parts to 31 in Kattegat in the southwest. It is relatively shallow in relation to its size, with the coastal zone constituting a large and important part of the ecosystem. Figure 1, from HELCOM (2010), illustrates the richness of habitat types (named ecosystem components) in different parts of the Baltic Sea. The categorization of the ecosystem components in this figure closely resembles the EFH categorization used in this review, apart from a few classes based on species data and deeper aphotic bottoms away from the coast, and can thus, in our opinion, be used as a proxy for EFH in the Baltic Sea.

In the context of Figure 1, an ecosystem component refers to biological parts of the ecosystem such as species, biotopes formed by habitat-forming species or abiotic biotopes with a clear linkage to certain species (Korpinen et al. 2012). The 14 named ecosystem components in Korpinen et al. (2012) are divided into benthic biotopes (two), benthic biotope complexes (six), water column (two) and species data (four). In the map the habitats specifically constitute: 1) mussel beds and 2) eelgrass meadows (benthic biotopes); 3) photic sand, 4) non-photoc sand, 5) photic mud and clay, 6) non-photoc mud and clay, 7) photic hard bottom and 8) non-photoc hard bottom (benthic biotope complexes); 9) photic water and 10) non-photoc water (water column); as well as 11) harbour porpoise, 12) seals, 13) seabird wintering grounds and 14) spawning and nursery areas of cod (species data). Note, however, that for the purposes of this review, a number of ecosystem components from the list above are not fully

synonymous to coastal EFH, as the term is interpreted and used in the present study. This clearly applies to the species data points 11–13 above, but also partly to non-photic bottoms (points 4, 6 and 8 above) and non-photic water column (point 10), i.e. for those parts that are occurring deeper down and farther away from the shoreline.



**Figure 1. Map showing the number of ecosystem components present (benthic and water column biotope complexes, benthic biotopes and species-related data layers) as a proxy for EFH in 5 km × 5 km squares in the Baltic Sea. Altogether 14 data layers were used when constructing the map, but no single square contained all ecosystem components. The map is taken from HELCOM (2010), with permission.**

If the ecosystem components from Korpinen et al. (2012) and coastal EFH in the Baltic Sea are considered to be of the same kind, the richest diversity of components/EFH is found in squares in the southwestern Baltic Sea, for example in the Sound, in the Belts and in Kattegat. A reasonably high diversity of components/EFH are also found around the large islands and in the archipelagos of the central Baltic Proper.

Lower diversities (fewer EFH) are found in the Bothnian Bay and in the eastern parts of the Baltic Sea (Figure 1).

## **2.2. Importance of coastal EFH in the Baltic Sea**

The importance of coastal EFH can in general be assessed as the effects of changes in their quantity or quality on metrics of viability and production of fish populations, stocks or communities in time or space (e.g. Levin and Stunz 2005, Sundblad et al. 2014). A recent review by Seitz et al. (2014) shows that in the Northeast Atlantic, 44% of all “ICES species”, i.e. species assessed and advised by the International Council for the Exploration of the Sea, utilizes coastal habitats as spawning, feeding, nursery or migration areas. These stocks contribute to 77% of the commercial landings of the “ICES species”. It follows then, that a limited habitat supply, possibly acting independently at different life-history stages utilising different habitats, can impact the size and dynamics of fish populations, although the relationships are not easily quantified (Seitz et al. 2014, Vasconcelos et al. 2014, Kallasvuori et al. 2017).

The available quantitative evidence for the importance of coastal habitats for fish production and viability has been achieved through a number of different approaches. These approaches include e.g. model based ones (e.g. Minns et al. 1996, Halpern et al. 2005, Levin and Stunz 2005, Fodrie et al. 2009), long-term field experiments (Schmitt and Holbrook 2000), otolith chemistry (e.g. Fodrie and Levin 2008), habitat specific biomass and size distributions (e.g. Mumby et al. 2004) and nursery habitat size (Rijnsdorp et al. 1992). Species distribution modelling has, in this respect, emerged as a promising tool to map specific habitat requirements for different life stages of species with ontogenetic habitat shifts (Bergström et al. 2013, Sundblad et al. 2014). By using modelling techniques, species occurrence or abundance can be related to map-based

predictor variables and thereby, fine-scale mapping of the distribution of species and habitats across spatially heterogeneous ecosystems can be carried out (Elith and Leathwick 2009, Pittman and Brown 2011, Bučas et al. 2013, Kotta et al. 2016, Moore et al. 2016).

### **2.2.1. Direct and indirect evidence of the effects of coastal EFH on fish population size**

From the Baltic Sea, some case studies give direct (quantitative) evidence on the role of coastal EFH for fish populations and fish production, although most of the evidence can be characterised as indirect (Table 1). Also, there do not seem to be any studies available from the Baltic Sea utilizing habitat-specific demographic rates, although this has been a preferred method for demonstrating habitat dependence in many circumstances globally (Levin and Stunz 2005, Vasconcelos et al. 2014). As may be noticed from the case studies below, the area of establishing direct links between habitats and fish populations is quite understudied in the Baltic Sea and most evidence seems to be available between habitats and larval fish, not directly for adult populations. Despite the fairly low number of studies showing direct links between fish stock sizes and availability of habitats, a reasonable amount of data on occurrence, or preferentially abundance, of various life stages of different fish species in specific habitats still indirectly indicate the importance of coastal EFH and help in their further identification and verification.

As direct evidence, Sundblad et al. (2014) used species distribution modelling on data from Sweden and Finland and related the distribution of nursery habitats for perch,

*Perca fluviatilis*, and pikeperch, *Sander lucioperca*, to the size of the adult populations of these species in twelve archipelago areas in the northern Baltic Proper. By doing this, the authors reveal that availability of coastal EFH explains almost half of the variation in population size, indicating a crucial role in limiting adult stock sizes. The relationships are, however, non-linear, suggesting that the negative effects of e.g. habitat loss or positive effects of e.g. restoration measures will be most significant in areas with the most limited habitat availability.

For whitefish, *Coregonus lavaretus*, Vanhatalo et al. (2012) utilized data from both the Swedish and Finnish coasts of the Gulf of Bothnia to establish direct relationships between environmental variables characterizing coastal EFH and larval production. Vanhatalo and colleagues used Gaussian processes for species distribution modelling and show that the most important variables describing potential larval areas over large scales, are bottom type, prolonged ice period in spring, ecological status of coastal areas, distance to large shallow sand areas and water depth. Thus, the most important variables are descriptors of coastal EFH for whitefish larvae and a metric of the current level of human impact on these areas.

In a recent Finnish case study, as a final example of direct connections between coastal EFH and coastal fish populations in the Baltic Sea, Kallasvuo et al. (2017) assessed the most important reproduction habitats for fish by using larval survey data and Bayesian species distribution models. By utilising data for four commercially and ecologically important fish species along the Finnish coast, Baltic herring (*Clupea harengus membras*), perch, pikeperch and smelt (*Osmerus eperlanus*), Kallasvuo and colleagues demonstrate that the production of fish stocks can be concentrated to very

limited areas compared to the total suitable production area that is available. Thus, spawning areas that are highly effective relative to the general pool of spawning areas can be identified. The applied methodology enables linking of the total production potential across the whole distribution area to fisheries stock assessment and management, especially for more strictly coastal species such as perch and pikeperch.

Concerning cod, *Gadus morhua*, there are a few studies available from the Baltic Sea that show the direct relationship between the volume of EFH for reproduction and the adult stock. MacKenzie et al. (2000) estimated reproduction volumes in time and space and demonstrate that the volume of EFH for egg survival determines the interannual stability in hatching success of cod eggs, while Cardinale and Arrhenius (2000) by the use of generalized additive models show that the volume of EFH for reproduction also affects cod recruitment. These results for cod are, however, not primarily focusing on coastal EFH. Still, with regard to coastal EFH, a recent study by Hinrichsen et al. (2017) demonstrates the importance of habitat availability for juvenile cod (nursery) and its effect on density-dependent growth, as a process relevant for recruitment success. Thus, across multiple life history stages, EFH availability influences stock size.

The remaining case studies presented in this chapter and in Table 1 are more indirect with regard to the connections between coastal EFH and fish populations, although there are no sharp distinctions between the direct studies mentioned above and the indirect ones mentioned below.

Hansen and Snickars (2014) utilized data from Sweden and Finland and report that the quality (species composition) of the macrophyte community on shallow soft bottoms

in relation to anthropogenic stressors shows good compliance with fish reproduction data. Bays that are dominated by stress sensitive macrophyte species also prove to be important nursery areas for fish. In another case study from Sweden and Finland, Snickars et al. (2010) report that distribution of spawning habitat for perch depends strongly on the type of substrate. The substrates generally consist of different types of vegetation, where the ones providing rigidity and structural complexity are preferred by the perch. Also, water depth, wave exposure and temperature matter to a relatively high extent with shallow depths and sheltered areas being preferred habitat characteristics. No direct links to the size of perch stocks have, however, been established.

In another case study from southern Finland, Engström-Öst et al. (2007) compared habitat choice and survival of pike larvae (*Esox lucius*) experimentally and conclude that pike larvae prefer and also survive better in filamentous algae (*Cladophora glomerata*) than in bladderwrack (*Fucus vesiculosus*) in the presence of predators. This is probably because the bladderwrack habitat is too “open” for the newly hatched pikes. In a related experimental study, Engström-Öst and Mattila (2008) compared the performance of larval pike under the influence of turbidity induced by phytoplankton. In this study, they report that the larval weight of pike is lower in turbid water, despite that pike larvae spend less time in vegetation and attack more prey. Thus, both direct (i.e. feeding and habitat choice) and indirect qualities (i.e. weight) of pike larvae are affected by the habitat quality (macroalgal structure, turbidity) and therefore probably also larval survival and recruitment to the adult population (Engström-Öst et al. 2007, Engström-Öst and Mattila 2008).



In a case study comprising the entire Finnish coastline, Uusitalo et al. (2012) used a Bayesian network model (expert driven model structure, data-learned parameters) to study the effects of many different factors (N, P, chlorophyll *a*, duration of ice coverage in winter, shore density in the area and salinity) on the CPUE (of reported commercial catches). Shore density was defined as the length of the shoreline within a rectangle, measured from the basic water level line from a 1:20 000 map and divided by the area of water surface in the rectangle (in ha), and it reflected the availability of coastal areas in the rectangle. The tested fish species were among others: pikeperch, pike, perch, flounder (*Platichthys flesus*), Baltic herring, burbot (*Lota lota*) and smelt. In their study, Uusitalo et al. (2012) report that shore density is the most influential factor. The strongest effects occur for pike, although it is concluded that shore density, corresponding closely to the availability of coastal EFH, is an important factor for all species, despite the fact that many of them are essentially freshwater ones, whose distribution also can be limited by salinity.

With regard to the importance of coastal EFH for production and viability of flounder, there are a number of case studies available from the Baltic Sea. In a study from Latvia, Ustups et al. (2013) utilized data spanning over 30 years to demonstrate that the spawning habitat (available water volume suitable for reproduction with regard to oxygen conditions) positively affects the survival and abundance of flounder larvae. Still, recruitment does not correlate with the supply of larvae, suggesting the presence of a bottleneck in the availability of juvenile growth habitat, which in itself, is also coastal. Case studies from southern Finland used fishery-independent data on adult flounder as well as historical and present-state data on juveniles in shallow coastal areas. These studies show that a pronounced decrease in abundance of juveniles correlates with an

increased bottom coverage of filamentous algae. A simultaneous decrease in the abundance of the adult stock indicates that a decline in the availability of EFH for juveniles acts as a bottleneck for the flounder population (Jokinen et al. 2015, 2016), supporting the conclusions of Ustups et al. (2013). Similar results have also previously been demonstrated by Pihl et al. (1994) and Carl et al. (2008) in the Kattegat and by Florin et al. (2009) for the Baltic Sea, but in the latter study more clearly for turbot (*Scophthalmus maximus*) than for flounder. The results for flounder above are further supported by Orio et al. (2017) who modelled spawning areas of flounder at a Baltic-wide scale and recognise a positive correlation between flounder spawning areas and adult stocks. The findings by Ustups et al. (2013) and Orio et al. (2017) are included as direct evidence in Table 1, although like the case with cod above, these results are not fully “coastal”.

For pikeperch in the German area of the Baltic Sea, the population size is strongly connected with the occurrence of suitable spawning sites in the inner coastal waters with lower salinities around 5-6 (Winkler 1996). These EFH are the base for nearly 40% of the total annual catch of pikeperch in German coastal waters with higher salinities (around 10) and corresponding numbers, or 44 %, can be shown for roach (Winkler et al. unpubl.).

For pike, Nilsson et al. (2014) show an increased recruitment of juveniles in three coastal wetlands of SE Sweden which have been restored in different ways. In areas with temporally flooded terrestrial vegetation, the migration of pike juveniles is shown to increase from a few thousand individuals in previous years to >100,000 individuals after the measures have been taken. To what extent these restored wetlands affect adult fish

stocks in coastal areas remains to be clarified, although there are indications of positive effects (Fredriksson et al. 2013).

Finally, some species utilize both coastal habitats, coastal wetlands and rivers for spawning and may display sympatric, genetically isolated populations. While their juvenile and adult stages may occur in the same habitats, spawning takes place in either fresh or brackish waters (Westin and Limburg 2002, Wastie 2014). The relative proportions of these sympatric populations may differ between areas and through time. In case studies from Estonia and Sweden, the relative importance of fresh or brackish water recruitment areas (spawning habitat preferences) for brackish water fish populations was examined through the use of otolith Sr:Ca profiles. These studies demonstrate the importance of coastal wetlands and rivers as spawning habitats for (semi-)anadromous fish as pike (Engstedt et al. 2010, Rohtla et al. 2012), burbot and ide (*Leuciscus idus*) (Rohtla et al. 2014, 2015). In the Väinameri Sea area in Estonia, 90% of adult pike hatches in fresh water and only 10% in brackish water (Rohtla et al. 2012; Rohtla 2015). In Sweden, 20% of pike hatches in brackish water in the Forsmark area at the 60° N latitude and 80% hatches in brackish water in the Kalmar Sound at the 56° N latitude (Engstedt et al. 2010). When compared with older (observational or anecdotal) data, the Estonian results suggest that brackish-water spawning pike is becoming rarer, which may be a result of deteriorated brackish water spawning grounds (Rohtla 2015). Along the Estonian coastal area, Rohtla et al. (2017) further demonstrate, also through the use of otolith chemistry techniques, that brackish water spawning whitefish has become rarer, which probably also reflects a poorer ecological status of its coastal spawning areas. Similarly, Byström et al. (2015) notice an important role of freshwater

habitats for perch recruitment in a Swedish coastal area with high abundance of the three-spined stickleback, which may prey on early life stage perch.

#### **2.2.2. Means to increase the knowledge of the importance of coastal EFH**

Although many different coastal habitats are essential for fish production and for the provisioning of rich fish communities in the Baltic Sea, the establishment of direct/quantitative relationships demonstrating their actual role for fish production is still in its infancy. The relatively low number of studies explicitly dealing with the importance of EFH for fish stocks is somewhat surprising. For many species, too little seems to be known about the ecology of the species in order to assess whether habitats are actually essential and limiting the production and viability of the populations (Levin and Stunz 2005, Seitz et al. 2014). Better evidence is, however, often found for non-migrating coastal species compared to migrating species (Iles and Beverton 2000), with cod (Hinrichsen et al. 2017) and the demersal ecotype of flounder (Orio et al. 2017) as possible exceptions. This could potentially be due to the conservative nature in habitat choice of non-migratory fish, or simply that it is easier to detect fish-habitat relationships in studies where many geographically restricted populations are included.

In other cases, indirect evidence exists or data for quantitative examination of the importance of coastal EFH for fish stocks may already be available and additional analyses could contribute to pinpoint their ecological importance (Pulkkinen et al. 2011, Kraufvelin et al. 2016). In a recent paper, Macura et al. (2016) present a methodological protocol for conducting a systematic review mainly on the impact of anthropogenic-induced physical and structural habitat changes on fish recruitment in shallow nearshore areas. Such a protocol can be used to assess the importance of undamaged coastal EFH for fish production. Further evidence on the role of coastal EFH can also be achieved using spatial approaches (e.g. assessing relationships between habitats of juveniles and adult fish to detect bottlenecks in early life stage), temporal data analyses (e.g. assessing variability between years in success of different life stages), stage-structured modelling (assessing habitat specific survival in stage-structured models) or otolith chemistry techniques (comparing contribution of different habitats through “fingerprinting” of different juvenile habitats). Currently, the most promising approach may be to estimate habitat-specific demographic rates in stage-structured modelling (Levin and Stunz 2005, Vasconcelos et al. 2014). It is then important, however, to combine this approach with habitat maps to quantify the importance of different habitats. When used properly, this approach may identify low productivity (per unit area) habitats as important, if they are abundant enough, compared to very productive habitats that are scarcer.

It should also be stressed that the establishment of a link between coastal EFH and fish stocks may not always be the prime interest as this is sustained already by the definition of EFH and the fact that a fish population is viable. Instead, the importance of EFH utilised by a population throughout different life history stages should maybe be

the centre of attention. This, in turn, leads to the question of “overlapping” EFH in a region or an area, and as a consequence, the difficulties to separate the relative effects or importance of different EFH (spawning, nursery, feeding, etc.) for a fish population and how “sub-EFH” are inter-linked and connected in the context of spatial/landscape ecology (Rose 2000, Levin and Stunz 2005, Vasconcelos et al. 2014).

### **3. THREATS TO AND PRESSURES ON COASTAL EFH IN THE BALTIC SEA**

#### **3.1. Generally about the conditions of coastal EFH in the Baltic Sea**

Coastal EFH in the Baltic Sea are exceptionally vulnerable as several natural features make the sea area inherently susceptible to the influence of human pressures. The Baltic Sea has a long water residence time (~30 years) and a large catchment area, which is relatively highly populated. The environmental status of many coastal areas of the Baltic Sea has declined considerably over the last 50 years (Bonsdorff et al. 1997, Lotze et al. 2006, Węśławski et al. 2013, Olsson et al. 2015, Andersen et al. 2015, Bergström et al. 2016a). This has for example led to evident changes in species composition of coastal fish, benthic invertebrate and macrophyte communities (e.g. Boström et al. 2002, Olsson et al. 2012, 2013, Snickars et al. 2015, Bergström et al. 2016a). The multifaceted environmental problems of the Baltic Sea, including extensive algal blooms, increasing areas of anoxic sea bottoms, contaminated organisms, and overexploitation of fish stocks, emerge as real challenges for environmental management calling for integrated strategies focusing on both fish and their preferred environments (e.g. Borja et al. 2016, Uusitalo et al. 2016). Within this process, a central focus on nearshore coastal areas


subjected to environmental pressure could be pivotal for the future potential of the Baltic Sea to provide ecosystem goods and services (Holmlund and Hammer 1999, Rönnbäck et al. 2007, Ahtiainen and Öhman 2014, Uusitalo et al. 2016).

As a spatial representation for weighing large numbers of cumulative anthropogenic impacts against ecosystem components and describing the current condition of various part of the sea area, the Baltic Sea Impact Index has been developed (see Halpern et al. 2008, HELCOM 2010, 2017 and table 2 in Korpinen et al. 2012 for details). This index shows that the lowest cumulative impact is generally found in the Gulf of Bothnia in the sparsely populated northernmost part of the Baltic Sea, and the highest impacts mainly occur in the coastal areas of the Finnish south and southwest, along the Estonian northern and western coast, along the east and west coast of southern Sweden, in the Polish Bay of Gdansk and in the Danish and German parts of the Baltic Sea (Figure 2). This impact map may be regarded as closely reflecting the general pressures on coastal EFH, as well.

Eutrophication, coastal construction and development, climate change, invasive species and fisheries have been acknowledged as major human-induced threats to coastal EFH in general (Jackson J.B.C. et al. 2001, Kappel 2005, Powers et al. 2005, Orth et al. 2006, HELCOM 2010, Hansen and Snickars 2014, Seitz et al. 2014, Sundblad and Bergström 2014, Kraufvelin et al. 2016). A specific feature and a natural threat to coastal EFH in the Baltic Sea is the post-glacial land-uplift process, which naturally, but constantly, shapes and alters the coastline and its shallow habitats for instance when semi-isolated flads and bays turn into freshwater ecosystems (Snickars et al. 2009, Meriste and Kirsimäe 2014). Among the human-induced threats, physical pressures

507 such as trawl fishery, shipping and boat traffic with the required infrastructure in the  
508 form of dredging, and shoreline modifications generally cause direct impacts on the  
509 habitats and are hence – in theory – easier to manage (Eriksson et al. 2004, Sandström et  
510 al. 2005, Sundblad and Bergström 2014, Pommer et al. 2016). Other (non-physical)  
511 threats/pressures usually act more indirectly and are hence often more challenging to  
512 manage (Elliott 2010, Duarte 2014). Most human-induced threats are severe on their  
513 own, but often have their largest impact when acting additively and synergistically  
514 (Elliott 2004, McLusky and Elliott 2004, Crain et al. 2008). Fish communities are affected  
515 both directly when exposed to these threats and indirectly through fragmentation,  
516 deterioration and loss of habitat. Here, the distinction between different fish species  
517 must again be stressed as for instance





**Figure 2. Presentation of cumulative potential anthropogenic impacts by the Baltic Sea Impact Index in 5 km × 5 km assessment units. The index in each assessment unit consists of the sum of anthropogenic impacts on selected ecosystem components present in the unit. The original index formula is from Halpern et al. (2008) and Korpinen et al. (2012). The map is taken from HELCOM (2017), with permission.**

mesopredatory fish, such as cyprinids and sticklebacks, may benefit from some of these threats/pressures or the negative effects of the threats/pressures imposed on other fish species (see e.g. Persson et al. 1991, Sandström and Karås 2002, Bergström et al. 2015, Byström et al. 2015). This may also be the case, to some extent, for pikeperch, which seems to be benefitting from coastal eutrophication and warmer summers (Heikinheimo et al. 2014) and also for the non-indigenous round goby *Neogobius melanostomus* (Ojaveer et al. 2015).

From a strict habitat perspective, there are some inherent differences with regard to which threats/pressures are the most dramatic ones for coastal EFH in the Baltic Sea. Seagrass and macrophyte beds are threatened by anthropogenic factors such as poor water quality caused by pollution, eutrophication, dredging, excessive sedimentation, altered openness of sheltered bays to the sea, climate change (leading to increased land runoff) and coastal development (Hemminga and Duarte 2000, Idestam-Almquist 2000, Airolidi and Beck 2007, Snickars et al. 2009, 2015, Rosqvist et al. 2010). Perennial macroalgal belts are threatened by eutrophication processes increasing the abundance of ephemeral algae, that suppress or inhibit the recolonization of canopy-forming algae and other organisms (Thompson et al. 2002, Råberg et al. 2005, Korpinen et al. 2007, Kraufvelin et al. 2007, 2010), but also by human construction and urbanization affecting water movement, water quality and causing habitat-related changes (Vogt and Schramm 1991, Eriksson et al. 1998, Kraufvelin 2007, Kraufvelin et al. 2010). Mussel beds are threatened by eutrophication, pollution, sedimentation, invasive species (e.g. the round goby), destructive fishing practices, and processes connected with climate change, such as higher water temperatures, acidification, increased storminess, increased land run-off and decreased salinity (Thompson et al. 2002, Airolidi and Beck 2007, Rakauskas et al.

2013, Díaz et al. 2015). Some of these pressures may, however, sometimes also prove to be beneficial, for instance for blue mussels (*Mytilus trossulus*) when new settlement areas are provided or when there are moderate increases in water movement (Díaz et al. 2015) and in temperature levels seasonally (Widdows 1991). The information on current threats to sedimentary environments, finally, is quite scarce (Brown and McLachlan 2002), but the major pressures on these habitats consist of the construction and use of marinas and ship ways including dredging, extraction of sand or gravel, trawl fishery, eutrophication, tourist developments, pollution from sewage discharge and industries as well as aquaculture activities (Newell et al. 1998, Airolidi and Beck 2007).

Thus, not all coastal EFH are affected by exactly the same threats, nor do they respond in the same way to similar pressures. All the human activities mentioned above are involved in causing different types of pressures and impacts on the habitats e.g. anoxic conditions in estuaries and enclosed basins (Karlson et al. 2002), accumulation of drifting algae (Vahteri et al. 2000), long-term accumulation of contaminants (Islam and Tanaka 2004) and introduction of non-indigenous species (Leppäkoski et al. 2002, Katsanevakis et al. 2014, Ojaveer and Kotta 2015). For more detailed information on species and habitats in the north-eastern Atlantic, see <http://www.marlin.ac.uk/>. Exclusively for the Baltic Sea, this kind of information is being gathered within HELCOM (<http://www.helcom.fi/>) and at a national level at least in Finland (<http://paikkatieto.ymparisto.fi/velmu/>).

### 3.2. Case studies about threats to and pressures on coastal EFH

Eutrophication favours the production of fast-growing, short-lived benthic and planktonic algae, that alters the structure and function of marine habitats and may cause hypoxia when accumulated and broken down (Lundberg 2005, Conley et al. 2009, Kraufvelin et al. 2010, Paerl and Otten 2013). This human pressure is acknowledged as a major problem to coastal EFH all over the Baltic Sea (HELCOM 2010, 2017, Kraufvelin et al. 2016). The large-scale decrease in distribution of the macroalga bladderwrack in eastern Sweden and southwestern Finland at the deeper end of its depth limit is of specific relevance for this study. This bladderwrack habitat loss is mainly caused by eutrophication-related processes in form of decreased light penetration and hampered recruitment and growth due to competition with filamentous algae and sedimentation (Kautsky et al. 1986, Korpinen et al. 2007, Kraufvelin et al. 2007, Rinne et al. 2011). As a consequence of this, large areas of shallow waters, potentially valuable for coastal fish, have been lost (Kautsky et al. 1986, Bergström et al. 2013, Vahteri and Vuorinen 2016). Similar patterns were also found in the shallow Puck Bay in Poland (Plinski and Florczyk 1984, Ciszewski et al. 1992, Węśławski et al. 2009), although this area is now slowly recovering (Węśławski et al. 2013). Another typical phenomenon due to eutrophication is the reed belt overgrowth of lagoons, sheltered bays and river mouths (Pitkänen et al. 2013, Altartouri et al. 2014, Meriste and Kirsimäe 2014). This process is potentially making shallow areas less useful as habitats for fish (Kneib and Wagner 1994, Weinstein and Balletto 1999), although see also Härmä et al. (2008), Lappalainen et al. (2008), Snickars et al. (2010) and Nilsson et al. (2014) for some positive influences of reed vegetation on fish communities, especially pike, but also for perch. Probably, too wide-spread and compact reed belts are negative for fish, while more restricted belts, and

belts from the previous season that have been flattened from ice and waves as well as the outer edges of reed areas are generally positive for fish (Lappalainen et al. 2008).

Eutrophication is also often acting in concert with other pressures such as coastal construction, seabed disturbance, climate change, overfishing and species introductions (Lundberg 2005) and understanding relationships between ecosystems and multiple human-induced pressures acting simultaneously is indeed a major challenge within marine environmental management (Borja 2014, Borja et al. 2016). Eutrophication combined with mesopredator release due to overfishing of large piscivorous fish species constitutes an example of a cumulative pressure, which can have strong effects on coastal EFH and present extensive challenges for management (Eriksson et al. 2009, 2011, Östman et al. 2016, Uusitalo et al. 2016). Eutrophication combined with the presence of invasive species can also impose interactive pressure on coastal EFH, as in the case with the recent invader in the northern Baltic Proper, Harris mud crab, *Rhithropanopeus harrisii*, occurring in both bladderwrack (Jormalainen et al. 2016) and eelgrass beds (Gagnon and Boström 2016) in the Finnish Archipelago Sea and in boulder fields with bladder-wrack (Nurkse et al. 2015) as well as in un-vegetated soft bottom areas in Estonia (Lokko et al. 2017). This invader acts as a mesopredator and can strongly reduce the number of grazers and impair their capability to buffer excessive growth of filamentous algae leading to decreased biodiversity and lowered habitat quality. Eutrophication effects combined with coastal construction dampening wave action can be exemplified by Kraufvelin et al. (2010) who conducted long-term experiments in outdoor rocky shore mesocosms. Kraufvelin and colleagues show that a combination of high nutrient enrichment with 50% lowered wave action over two years lead to a 2.5-fold reduction of habitat-forming perennial brown algae (mainly of the

order Fucales) and an 80-fold increase in annual green algae (mainly of the order Ulvales).

The physical pressure from human activities is both high and increasing in the coastal zone, especially in the shallowest areas and habitats (Sundblad and Bergström 2014). Activities such as recreational boating, building of marinas and other forms of construction constitute major problems for coastal EFH all over the Baltic Sea, but perhaps currently to a higher extent in Sweden, Finland, Poland, Germany and Denmark than in Estonia, Latvia and Lithuania (HELCOM 2010, Dafforn et al. 2015, Kraufvelin et al. 2016). In the Stockholm archipelago of Sweden, Sundblad and Bergström (2014) used predictive habitat modelling and mapping of human pressures to estimate the cumulative long-term effects of coastal development in relation to fish habitats. The results suggest an annual increase in the proportion of degraded areas of 0.5% on average and of 1% for areas close to larger human population centres. Furthermore, the same study shows that approximately 40% of available habitat for pike, perch and roach was already subject to some form of construction by 2005 (Sundblad and Bergström 2014).

In Estonia, Latvia, Lithuania and Poland, invasive species are, apart from eutrophication, brought forward as important human-induced threats to coastal EFH (HELCOM 2010, Kraufvelin et al. 2016). Among invasive species, the round goby has been of increasing importance during the last years (Ojaveer et al. 2015, Kotta et al. 2016) with potential to impact the distribution of EFH in the form of blue mussel beds (Järv et al. 2011, Kornis et al. 2012, Rakauskas et al. 2013). Round gobies generally prefer hard bottom habitats, where mussels make up its most important food source

(Barton et al. 2006, Karlson et al. 2007, Järv et al. 2011, Kornis et al. 2012, Rakauskas et al. 2013), although Nurkse et al. (2016) characterize the species as a generalist consumer. Due to competition with round gobies, it has also been shown that juvenile turbot change their diet and turbot recruitment simultaneously decreases significantly (Ustups et al. 2016). Round gobies may also, through competition for food and habitat, negatively affect flounder (Karlson et al. 2007, Järv et al. 2011, Orio et al. 2017), ruffe *Gymnocephalus cernua* (Rakauskas et al. 2013), and viviparous eelpout *Zoarces viviparus* (<https://www.nobanis.org/marine-identification-key/fish/fish-start/fish-key/neogobius-melanostomus/>). The effects of invasive species increase as the populations establish and spread to adjacent areas as can be seen with the round goby in the southwestern Baltic Sea (Azour et al. 2015). The round goby may, however, not only influence the biological communities of the Baltic Sea negatively. Recent studies from the northeastern German coast (Oesterwind et al. 2017) and from Estonia (Liversage et al. 2017) show that the round goby is included in the local food web, including fish eating birds.

In Germany, Denmark and on the southern and southwestern coast of Sweden, major human-induced threats to coastal EFH are, in addition to eutrophication and climate change, coastal construction, demersal trawling, tourism, dredging and material extraction (HELCOM 2010, Kraufvelin et al. 2016). Material extraction, e.g. extensive removal of stones and boulders in coastal areas of Denmark has not only led to destruction of reefs and removal of hard bottom habitat, but also to the loss of biogenic structures associated with and characteristic of these reefs (Carr 1994, Dahl et al. 2008). Støttrup et al. (2014) studied a re-established stony reef in Kattegat and documented an increase in fish abundance and can thereby demonstrate that these damages may be to



some extent reversible. Also, bottom trawling in the Kattegat has led to a decrease in hard bottoms in general through removal of stones and boulders (homogenisation of mixed bottoms) and to a decrease in the amount of sensitive species, some of which are habitat-forming (Hopkins 2003, Pommer et al. 2016).

Interestingly, despite many scientists mentioning climate change as a major threat to coastal EFH in their regions (Kraufvelin et al. 2016), there are still few studies from the Baltic Sea that explicitly focus on climate change related effects on EFH. This is surprising as many different pressures in the Baltic Sea fall under the climate change umbrella such as increased temperatures, decreased salinity, decreased oxygen concentrations, acidification, increased storminess, increased sea levels, etc. (BACC Author Team 2008, HELCOM 2013). There are, however, some references available that are related to effects on coastal EFH, e.g. for macrophytes from the Baltic Proper (Idestam-Almquist 2000, Härmä et al. 2008), for perennial bladderwrack from the Baltic Proper and from the southwestern Baltic Sea (Kraufvelin et al. 2012, Graiff et al. 2015, 2017), for blue mussels from the southwestern Baltic Sea (Thomsen et al. 2010, Havenhand 2012), and for fish and zoobenthos from the entire Baltic Sea (MacKenzie et al. 2007) and from the Baltic Proper (Snickars et al. 2015), although most of the reported and projected habitat effects in these studies are rather minor ones.

To better quantify and evaluate the magnitude of all threats to and pressures on coastal EFH highlighted in the case studies above and to provide more accurate and reliable information and recommendations for the management and conservation of EFH in a Baltic Sea wide perspective, maps of pressure variables, together with a mechanistic understanding of habitat effects of different threats/pressures, need to be



integrated with habitat maps. For these kinds of purposes, web-based knowledge platforms such as the one developed by MarLin for the UK (<http://www.marlin.ac.uk/>) can be utilized and applied. An attempt in this direction has also been done by HELCOM (2010) and Korpinen et al. (2012), as may be seen in Figure 2 of this review. More recent web resources can be found in HELCOM HOLAS II (<http://helcom.fi/helcom-at-work/projects/holas-ii/>, see also HELCOM 2017) and the associated HELCOM TAPAS (<http://helcom.fi/helcom-at-work/projects/tapas>). A promising approach to assess habitat quality based on the ecological status of benthic indicators and the EU Habitats Directive (Anon. 1992) has also recently been presented for Estonian waters by Torn et al. (2017) and similar approaches could be further developed for other regions of the Baltic Sea. Another way forward could be to combine probabilistic Bayesian network models describing the complex relationships between human activities and sensitive ecosystem components (e.g. sensitive habitats), with GIS databases (Stelzenmüller et al. 2010, Helle et al. 2016).

#### **4. INTEGRATED MANAGEMENT AND CONSERVATION OF COASTAL EFH IN THE BALTIC SEA**

The increasing anthropogenic impacts on marine waters have fuelled the discussion on how to manage and to conserve marine resources sustainably. During the last decade, there has been a raised focus on ecosystem-based management of marine ecosystems to secure the maintenance of healthy, productive, and resilient ecosystems capable of providing the services needed for the well-being of society (Collie et al. 2013, Yáñez-Arancibia et al. 2013, Borja 2014, Andersen and Kallenbach 2016, Borja et al. 2016). Within the Baltic Sea region, the current leading directives and agreements for this are the EU Marine Strategy Framework Directive (MSFD; Anon. 2008), the HELCOM Baltic Sea Action Plan (BSAP; HELCOM 2007) and the Common Fisheries Policy (CFP; Anon. 2013), but also the EU Habitats Directive (HD; Anon. 1992), the EU Water Framework Directive (WFD; Anon. 2000) and the EU Maritime Spatial Planning Directive (MSPD; Anon. 2014) are important.

Although both healthy fish populations and benthic habitats are central elements for maintaining a good status of the coastal environment, management of fisheries and nature conservation have historically been separated in the Baltic Sea region like in many other parts of the world (Sissenwine and Symes 2007, Kenny et al. 2009, Kraufvelin et al. 2016). The awareness of potential synergies between the two has also been low. Traditional management of marine resources has typically ignored interactions between fisheries and the status of coastal habitats, cross-system fluxes,

predator-prey interactions and other ecosystem components. An ecosystem-based management perspective where conservation and fisheries issues are integrated could instead provide mutual benefits and has therefore been brought forward as a more convenient platform for coastal systems (Pikitch et al. 2004, Leathwick et al. 2008, Thrush and Dayton 2010, Möllmann et al. 2014). Such a perspective would better cover the traits and needs of whole ecosystems and not only the ones of certain species, while simultaneously ensuring that multidisciplinary scientific approaches are adopted and that the right actors and stakeholders are involved (Hopkins et al. 2011, Long et al. 2015). The multitude of drivers to account for, however, also calls for other management strategies. With regard to the management of threats/pressures, cumulative impact assessments could be a functional approach for setting limits on allowable levels of human impact on ecosystems (Halpern et al. 2008, Korpinen et al. 2012, Rahikainen et al. 2014, Andersen et al. 2015, HELCOM 2017).

The conservation of coastal EFH is generally poor in the Baltic Sea, although coastal benthic habitats, and thus EFH, in many countries around the Baltic Sea, have been a focus of national conservation efforts (Sundblad et al. 2011). Within the fisheries management sector, attention has, however, mainly been devoted to commercial and threatened species. Maintenance and restoration of fish stocks have indeed been objectives in nature conservation, but still with restricted focus on the habitats themselves and with most focus directed towards salmonids or the threatened species covered by the Habitats Directive (Anon. 1992) and the European and national IUCN red-lists ([http://ec.europa.eu/environment/nature/conservation/species/redlist/index\\_en.htm](http://ec.europa.eu/environment/nature/conservation/species/redlist/index_en.htm),

Kraufvelin et al. 2016). Sundblad et al. (2011) investigated the representativity and connectivity of Marine Protected Area (MPA) networks in the northern Baltic Proper (Sweden and Finland) with respect to a coastal fish assemblage and associated habitats based on fish distribution maps and the linking of specific life stage occurrences to environmental variables. These analyses reveal that both the representativity and the connectivity of the network are poor as only 3.5% of the assemblage recruitment habitat is protected and 48% of potentially connected habitats are included in the MPA network. Furthermore, from a coastal EFH perspective, it appears that the most relevant areas are not always the ones being preserved. The lack of an ecosystem-based management perspective and the traditional split of fisheries and environmental management have again been major reasons underlying the poor conservation status of EFH in the Baltic Sea. Further challenges to the management of fish and habitats in the coastal zone are that they are under national jurisdictions of ten different countries in the Baltic Sea area, which cause large practical differences in management regimes. Hence, the authors see a need for the EFH perspective to be more strongly considered at both national and international levels of coastal management and conservation. Currently, changes appear to be taking place in many countries around the Baltic Sea (Kraufvelin et al. 2016) so now would be an opportune time for science advisors to bring EFH to the forefront of policy makers' attention.

In order to aid in merging the management of fisheries and environmental issues, there is a general need for a common awareness of the importance of coastal EFH and also about the threats to these habitats among managers, politicians and the public (e.g. Lotze 2004). There has been an apparent lack of information on the importance of the habitats for fish production and viability, but also previously a lack of maps depicting

the spatial distribution of specific types of coastal EFH to be used in marine spatial planning, permitting processes and for other management purposes. To that end, there is also a great need for more species- and life-stage-specific knowledge, both in terms of population-level effects and the geographical distribution of coastal EFH. As quantitative evidence for habitat limitation of fish production is slowly accumulating from different areas and species through the use of various methods (Vasconcelos et al. 2014, Seitz et al. 2014), the possibilities for integrating fish habitats in fisheries management and nature conservation will improve accordingly.

In order to reach a higher level of protection of coastal EFH, the role of habitats in supporting fisheries must also be disentangled in a broader context so that the value of the ecosystem services that these habitats provide can be emphasized more strongly (Holmlund and Hammer 1999, Rönnbäck et al. 2007, Uusitalo et al. 2016). These services may include producing fish for commercial and recreational fisheries, aquaculture and biological regulation (e.g. regulation of eutrophication symptoms through top-down control of filamentous algae), but also maintenance of biodiversity and ecosystem resilience. Many habitats considered EFH are also of importance for coastal protection against erosion, as nutrient filters, carbon sinks and for human recreation and scientific, educational and cultural purposes (Ahtiainen and Öhman 2014, Bouma et al. 2014, Ivarsson et al. 2017). Natural scientists together with environmental economists and social scientists should therefore consider all the ways in which coastal fish habitats provide value to society and use these as examples in communicating the needs for protection of coastal EFH and their sustainability (Støttrup et al. 2016). In this context, the general protection of coastal EFH from diverse pressures and what level of sustainable use that can be permitted should also be clearly stated

(Turner et al. 1999, Fluharty 2000). This information can be included in utility functions of decision support tools and in that way be accounted for when the performance of alternative management strategies/actions are evaluated quantitatively (see Laurila-Pant et al. 2015).

In the process of developing an efficient management scheme for the protection of coastal EFH, merging the objectives of fisheries and environmental management, possible difficulties of a common management of habitats and fish must be taken into careful consideration (Rose 2000, Rice 2005). Most exploited fish are long-lived, utilise many different habitats during their life cycle, and often exhibit large fluctuations in abundance. Efficient management therefore requires understanding how environmental variability, due to both natural and anthropogenic sources, affects fish population dynamics. Rose (2000) described a number of issues that are related to quantifying effects of environmental quality on fish populations and which at the same time may serve as demonstrations of how modelling could be used to address them. These issues include difficulties with the detectability of relationships, uncertainties due to heterogeneity in the habitat and disproportional population responses, unnecessary sacrifice of biological realism, neglected significance of community interactions, and ignored sublethal and cumulative effects. The quantification of effects of environmental quality on fish populations can be improved if these issues are carefully considered in the analyses, and by adopting multidisciplinary approaches that combine stage-structured modelling and life history theory (Rose 2000, Levin and Stunz 2005, Vasconcelos et al. 2014).

Finally, the need to combine alternative management strategies or actions and the objectives of the society, i.e. decision-making criteria against which success or failure of management are to be evaluated, should also be explored. Laurila-Pant et al. (2015) discuss these issues in connection with criteria setting towards a more holistic framework, although mainly with focus on biodiversity-related objectives. From a risk management perspective, approaches based on the precautionary principle may also sometimes be needed (Long et al. 2015, Chapman 2017). This, because uncertainty and lack of sufficient evidence are not acceptable reasons for not protecting supposedly essential habitats, if losing them may cause the collapse of fish stocks, with effects potentially propagating throughout food webs. Another issue which may complicate joint management is the inconsistency in the definition and understanding of the term habitat and habitat-related concepts in general (Elliott S.A.M. et al. 2016). It will not be dwelled further into habitat definitions in this review, but according to Elliott S.A.M. et al. (2016), unclear use of habitat-related terminology could have implications for the effectiveness of ecosystem-based fisheries management when e.g. different actors within marine science use the same terms with different connotations. However, when coastal management is implemented at local scales, the inclusion of all stakeholders from an early stage can go some way to mitigate such incommensurable language barriers and potential miscommunication (Hopkins et al. 2011).

## 5. SYNTHESIS

Coastal EFH form elementary cornerstones of the Baltic Sea ecosystem due to the central importance of fish for ecosystem functioning and the dependence of fishes upon specific coastal habitat types. As such, there are strong needs to focus on the protection of coastal EFH in addition to increasing our understanding of their species-specific importance, and on disentangling causal factors, pressures and mechanisms behind the changes that are observed in their status. Efficient management measures can be developed based on improved knowledge of causal factors and mechanisms for ecosystem change, e.g. how various stressors interact to structure communities (e.g. Rose 2000). The same applies to monitoring, assessing and mapping the availability and the state of coastal EFH as well as the documentation of human activities and pressure variables related to them (HELCOM 2010, Kraufvelin et al. 2016). Initial steps to bring this work forward could be to construct roadmaps, focus on directed studies and develop and harmonize the methodology (Kraufvelin et al. 2016). During this process, there will be evident needs for intensified cooperation between the Baltic Sea countries in order to reach successful implementations of international agreements and legislative marine acts such as the Baltic Sea Action Plan (BSAP), the Habitat Directive (HD), the



Marine Strategy Framework Directive (MSFD) and the Marine Spatial Planning Directive (MSPD). At the same time, local/regional conditions and the actual characteristics of the targeted ecosystems need to be taken into consideration more efficiently, because, as it has been shown in a number of cases in this review, the most efficient management may sometimes benefit from being planned and implemented case-specifically.

A major underlying objective for developing a more efficient management framework should be to improve the possibilities for connecting fisheries and environmental management across sectors (Pikitch et al. 2004, Thrush and Dayton 2010). Efforts made in these directions will also simultaneously aid in improving the sustainability of coastal EFH, enhance our abilities to predict and mitigate current and future effects of environmental change as well as support activities to create and implement adaptive management plans. To increase the awareness of the benefits of integrating management of fish and habitats, the scientific community can contribute in many ways. Ecological synergies achieved by protecting coastal EFH can be demonstrated; methods for large-scale mapping of EFH can be developed and utilized; effects of different threats to EFH may be quantified; and the importance of the habitats may be communicated (Kraufvelin et al. 2016). However, since not all habitats can be conserved or restored, some general frameworks to prioritize critical habitats of e.g. exploited fishes or red-listed species need to be developed and followed (Rose 2000). It must also be kept in mind that if a specific fish habitat is not strictly limiting population growth, a change in its availability does not lead to a change in stock sizes, provided that other regulating factors remain constant (Levin and Stunz 2005, Rice 2005).

This review gives an overview of the current knowledge as well as the lack of knowledge about coastal EFH in the Baltic Sea and brings about some suggestions for future work and cooperation. The topic is timely and of high importance in the current era of rapidly improving habitat modelling, new demands for better monitoring of marine ecosystem such as BSAP (HELCOM 2007), MSFD (Anon. 2008) and MSPD (Anon. 2014), and the findings that the Baltic network of MPAs cannot be considered ecologically coherent (Sundblad et al. 2011). The review also stresses the importance to protect key habitats vital for the survival of early life stages of fish and to map these areas (Kraufvelin et al. 2016). Apart from the need for conducting more investigations into the topics mentioned above, further studies also seem to be especially urgent within the field of attaining quantitative data for the value of coastal EFH for fish production, including defining the key habitats for protection and for possible restoration efforts, as well as disentangling the major threats/pressures and their effects (e.g. Elliott 2004, Elliott, M. et al. 2016). Improved integration of habitat quality in fish stock assessment and ecosystem-based fishery management is also warranted when this path is followed (Seitz et al. 2014, Sundblad et al. 2014). A crucial part of this work could consist of carrying out additional analyses on existing data as a lot of the needed information already seems to be available through monitoring and mapping work carried out in Baltic Sea countries (Kraufvelin et al. 2016). During this process, the utilization of meta-analytical approaches could be worth considering (see e.g. Pulkkinen et al. 2011, Östman et al. 2016). The initiation of common research projects and intensified outreach efforts constitute fruitful ways to bring this work forward on a Baltic-wide scale. In order to succeed with all these undertakings, devoted endeavours focusing on all aspects of coastal EFH will be of utmost importance. Successful implementation of these activities

will then in turn hopefully lead to clear and lasting improvements for fish and their habitats in the entire Baltic Sea region.

## 6. ACKNOWLEDGEMENTS

This review has been prepared on the basis of information collected and discussed during a workshop, 'Essential Coastal Habitats for Fish', in Öregrund, Sweden during June 2<sup>nd</sup>–4<sup>th</sup> 2015, gathering together 30 national experts from eight Baltic Sea countries. The workshop and preparation of the review article was funded by the Nordic Council of Ministers to Jens Olsson at Swedish University of Agricultural Sciences. The funding source had no practical involvement in the work carried out for the review article.

## 7. REFERENCES

Ahtiainen, H., Öhman, M.C., 2014. Ecosystem services in the Baltic Sea: valuation of marine and coastal ecosystem services in the Baltic Sea. *Tema Nord* 2014:563. Nordic Council of Ministers, Copenhagen, 74 pp.

Airolidi, L., Beck, M.W., 2007. Loss, status and trends for coastal marine habitats of Europe. *Oceanography and Marine Biology: an Annual Review* 45, 345-405.

Altartouri, A., Nurminen, L., Jolma, A., 2014. Modeling the role of the close-range effect and environmental variables in the occurrence and spread of *Phragmites australis* in four sites on the Finnish coast of the Gulf of Finland and the Archipelago Sea. *Ecology and Evolution* 4, 987-1005.

Andersen, J.H., Kallenbach, E., 2016. A literature study of human activities and pressures as well as ecosystem component layers available for Marine Spatial Planning and mapping of cumulative impacts in Swedish marine waters. NIVA Denmark Water Research 6997-2016. 36 pp.

Andersen, J.H., Halpern, B.S., Korpinen, S., Murray, C., Reker, J., 2015. Baltic Sea biodiversity status vs. cumulative human pressures. *Estuarine, Coastal and Shelf Science* 161, 88-92.

Anon., 1992, Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Official Journal of the European Union* 206, 7-50.

Anon., 2000. Directive 2000/60/EC of the European Parliament and of the council of 23 October 2000 establishing a framework for Community action in the field of water policy. *Official Journal of the European Union* L327, 1-72.

Anon., 2008. Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy (Marine Strategy Framework Directive). *Official Journal of the European Communities* L 164/19.

Anon., 2013. Regulation (EU) No 1380/2013 of the European parliament and of the council of 11 December 2013 on the Common Fisheries Policy, amending Council Regulations (EC) No 1954/2003 and (EC) No 1224/2009 and repealing Council Regulations (EC) No 2371/2002 and (EC) No 639/2004 and Council Decision 2004/585/EC.

Anon., 2014. Directive 2014/89/EU of the European Parliament and of the Council of 23 July 2014 establishing a framework for maritime spatial planning. *Official Journal of the European Union* L 257/135.

Armstrong, C.W., Falk-Petersen, J., 2008. Habitat-fisheries interactions: a missing link? *ICES Journal of Marine Science* 65, 817-821.

Aro, E., 1989, A review of fish migration patterns in the Baltic. *Rapports et Procès-verbaux des Réunions. Conseil International pour l'Exploration de la Mer* 190, 72-96.

- 1021 Azour, F., van Deurs, M., Behrens, J., Carl, H., Hüssy, K., Greisen, K., Ebert, R., Møller,  
1022 P.R., 2015. Invasion rate and population characteristics of the round goby *Neogobius*  
1023 *melanostomus*: Effects of density and invasion history. *Aquatic Biology* 24, 41–52.
- 1024 BACC Author Team, 2008. Assessment of climate change for the Baltic Sea basin.  
1025 Springer Science & Business Media.
- 1026 Barton, D.R., Johnson, R.A., Campbell, L., Petruniak, J., Patterson, M., 2005. Effects of  
1027 round gobies (*Neogobius melanostomus*) on dreissenid mussels and other invertebrates  
1028 in eastern Lake Erie, 2002–2004. *Journal of Great Lakes Research* 31, 252–261.
- 1029 Beck, M.W., Heck, K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M.,  
1030 Halpern, B., Hays, C.G., Hoshino, K., Minello, T., Orth, R.J., Sheridan, P.F., Weinstein, M.P.,  
1031 2001. The identification, conservation, and management of estuarine and marine  
1032 nurseries for fish and invertebrates. *Bioscience* 51, 633–641.
- 1033 Benaka, L.R. (Ed.), 1999. Fish habitat: Essential fish habitat and rehabilitation.  
1034 American Fisheries Society Symposium 22, Bethesda, MD.
- 1035 Bergström, L., Heikinheimo, O., Svirgsden, R., Kruze, E., Ložys, L., Lappalainen, A., Saks,  
1036 L., Minde, A., Dainys, J., Jakubavičiūtė, E., Ådjers, K., Olsson, J., 2016a. Long term changes  
1037 in the status of coastal fish in the Baltic Sea. *Estuarine, Coastal and Shelf Science* 169, 74–  
1038 84.
- 1039 Bergström, L., Bergström, U., Olsson, J., Carstensen, J. 2016b. Coastal fish indicators  
1040 response to natural and anthropogenic drivers–variability at temporal and different  
1041 spatial scales. *Estuarine, Coastal and Shelf Science* 183, 62–72.
- 1042 Bergström, L., Karlsson, M., Bergström, U., Pihl, L., Kraufvelin, P., 2016c. Distribution  
1043 of mesopredatory fish determined by habitat variables in a predator-depleted coastal  
1044 system. *Marine Biology* 163, 201.

- 1045 Bergström, U., Sundblad, G., Downie, A.-L., Snickars, M., Boström, C., Lindegarth, M.,  
1046 2013. Evaluating eutrophication management scenarios in the Baltic Sea using species  
1047 distribution modelling. *Journal of Applied Ecology* 50, 680-690.
- 1048 Bergström, U., Olsson, J., Casini, M., Eriksson, B. K., Fredriksson, R., Wennhage, H.,  
1049 Appelberg, M., 2015. Stickleback increase in the Baltic Sea – A thorny issue for coastal  
1050 predatory fish. *Estuarine, Coastal and Shelf Science* 163, 134-142.
- 1051 Blaber, S.J.M., Blaber, T.G., 1980. Factors affecting the distribution of juvenile  
1052 estuarine and inshore fish. *Journal of Fish Biology* 17, 143-162.
- 1053 Bonsdorff, E., Pearson, T.H., 1999. Variation in the sublittoral macrozoobenthos of the  
1054 Baltic Sea along environmental gradients: A functional-group approach. *Australian*  
1055 *Journal of Ecology* 24, 312-326.
- 1056 Bonsdorff, E., Blomqvist, E.M., Mattila, J., Norkko, A., 1997. Coastal eutrophication:  
1057 causes, consequences and perspectives in the archipelago areas of the northern Baltic  
1058 Sea. *Estuarine, Coastal Shelf Science* 44, 63-72.
- 1059 Borja, A., 2014. Grand challenges in marine ecosystems ecology. *Front Mar Sci* 1:1.
- 1060 Borja, A., Elliott, M., Andersen, J.H., Berg, T., Carstensen, J., Halpern, B.S., Heiskanen,  
1061 A.S., Korpinen, S., Lowndes, J.S.S., Martin, G., Rodriguez-Ezpeleta, N., 2016. Overview of  
1062 Integrative Assessment of Marine Systems: The Ecosystem Approach in Practice.  
1063 *Frontiers in Marine Science* 3, 20.
- 1064 Boström, C., Bonsdorff, E., Kangas, P., Norkko, A., 2002. Long-term changes of a  
1065 brackish-water eelgrass (*Zostera marina* L.) community indicate effects of coastal  
1066 eutrophication. *Estuarine, Coastal and Shelf Science* 55, 795-804.
- 1067 Bouma, T.J., van Belzen, J., Balke, T., Zhu, Z., Airoidi, L., Blight, A.J., Davies, A.J., Galvan,  
1068 C., Hawkins, S.J., Hoggart, S.P.G., Lara, J.L., Losada, I.J., Maza, M., Ondiviela, B., Skov, M.W.,  
1069 Strain, E.M., Thompson, R.C., Yang, S., Zanuttigh, B., Zhang, L., Herman, P.M.J., 2014.

Identifying knowledge gaps hampering application of intertidal habitats in coastal protection: Opportunities & steps to take. *Coastal Engineering* 87, 147-157.

Brown, A.C., McLachlan, A., 2002. Sandy shore ecosystems and the threats facing them: some predictions for the year 2025. *Environmental Conservation* 29, 62-77.

Bučas, M., Bergström, U., Downie, A.L., Sundblad, G., Gullström, M., von Numers, M., Šiaulys, A., Lindegarth, M., 2013. Empirical modelling of benthic species distribution, abundance, and diversity in the Baltic Sea: evaluating the scope for predictive mapping using different modelling approaches. *ICES Journal of Marine Science* 70, 1233-1243.

Byström, P., Bergström, U., Hjalten, A., Ståhl, S., Jonsson, D., Olsson, J., 2015. Declining coastal piscivore populations in the Baltic Sea: Where and when do sticklebacks matter? *Ambio* 44, 462-471.

Cardinale, M., Arrhenius, F., 2000. The influence of stock structure and environmental conditions on the recruitment process of Baltic cod estimated using a generalized additive model. *Canadian Journal of Fisheries and Aquatic Sciences* 57, 2402-2409.

Carl, J., Sparrevohn, C., Nicolajsen, N., Støttrup, J.G. 2008. Substratum selection by juvenile flounder *Platichthys flesus* (L.): effect of ephemeral filamentous macroalgae. *J. Fish Biology* 72, 2570-2578.

Carr, M.H., 1994. Effects of macroalgal dynamics on recruitment of a temperate reef fish. *Ecology* 75, 1320-1333.

Cattirjse, A., Hampel, H., 2006. European intertidal marshes: a review of their habitat functioning and value for aquatic organisms. *Marine Ecology Progress Series* 324, 293-307.

Chapman, P.M., 2017. Assessing and managing stressors in a changing marine environment. *Marine Pollution Bulletin* 124, 587-590.



- 1094 Ciszewski, P., Kruk-Dowgiallo, L., Zmudzinski, L., 1992. Deterioration of the Puck Bay  
1095 and biotechnical approaches to its reclamation. In: Bjørnstad, E., L. Hagerman, K. Jensen  
1096 (Eds.), Proc. 12th Baltic Marine Biologists symp. Olsen & Olsen, Fredensborg, pp. 43-46.
- 1097 Collie, J.S., Adamowicz, W.L., Beck, M.W., Craig, B., Essington, T.E., Fluharty, D., Rice, J.,  
1098 Sanchirico, J.N., 2013. Marine spatial planning in practice. *Estuarine Coastal and Shelf*  
1099 *Science* 117, 1-11.
- 1100 Conley, D.J., Björck, S., Bonsdorff, E., Carstensen, J., Destouni, G., Gustafsson, B. G.,  
1101 Hietanen, S., Kortekaas, M., Kuosa, H., Meier, H.E.M., Müller-Karulis, B., Nordberg, K.,  
1102 Norkko, A., Nürnberg, G., Pitkänen, H., Rabalais, N., Rosenberg, R., Savchuk, O.P., Slomp,  
1103 C.P., Voss, M., Wulff, F., Zillén, L., 2009. Hypoxia-related processes in the Baltic Sea.  
1104 *Environmental Science and Technology* 43, 3412-3420.
- 1105 Crain, C.M., Kroeker, K., Halpern, B.S., 2008. Interactive and cumulative effects of  
1106 multiple human stressors in marine systems. *Ecology Letters* 11, 1304-1315.
- 1107 Dafforn, K.A., Mayer-Pinto, M., Morris, R.L., Waltham, N.J., 2015. Application of  
1108 management tools to integrate ecological principles with the design of marine  
1109 infrastructure. *Journal of Environmental Management* 158, 61-73.
- 1110 Dahl, K., Stenberg, C., Lundsteen, S., Støttrup, J., Dolmer, P., Tendal, O.S., 2008. Ecology  
1111 of Læsø Trindel - A reef impacted by extraction of boulders. Danish Forest and Nature  
1112 Agency, 54 pp.
- 1113 Díaz, E.R., Erlandsson, J., Westerborn, M., Kraufvelin, P., 2015. Depth-related spatial  
1114 patterns of sublittoral blue mussel beds and their associated macrofauna diversity  
1115 revealed by geostatistical analyses. *Marine Ecology Progress Series* 540, 121-134.
- 1116 Duarte, C.M., 2014. Global change and the future ocean: a grand challenge for marine  
1117 sciences. *Frontiers in Marine Science*, 1, 63.

- 1118 Eddy, T.D., Lotze, H.K., Fulton, E.A., Coll, M., Ainsworth, C.H., de Araújo, J.N., Bulman,  
1119 C.M., Bundy, A., Christensen, V., Field, J.C., Gribble, N.A., Hasan, M., Mackinson, S.,  
1120 Townsend, H., 2017. Ecosystem effects of invertebrate fisheries. *Fish and Fisheries*, 18:  
1121 40–53.
- 1122 Elith, J., Leathwick, J.R., 2009. Species distribution models: ecological explanation and  
1123 prediction across space and time. *Annual Review of Ecology, Evolution and Systematics*  
1124 40, 677-697.
- 1125 Elliott, M., 2004. Marine habitats: loss and gain, mitigation and compensation. *Marine*  
1126 *Pollution Bulletin* 49, 671-674.
- 1127 Elliott, M., 2010. Marine science and management means tackling exogenic  
1128 unmanaged pressures and endogenic managed pressures – a numbered guide. *Marine*  
1129 *Pollution Bulletin* 62, 651-655.
- 1130 Elliott, M., Mander, L., Mazik, K., Simenstad, C., Valesini, F., Whitfield, A., Wolanski, E.,  
1131 2016. Ecoengineering with Ecohydrology: Successes and failures in estuarine  
1132 restoration. *Estuarine Coastal and Shelf Science* 176, 12-35.
- 1133 Elliott, S.A.M., Milligan, R.J., Heath, M.R., Turrell, W.R., Bailey, D.B., 2016. Disentangling  
1134 habitat concepts for demersal marine fish management. *Oceanography and Marine*  
1135 *Biology: An Annual Review* 54, 173-191.
- 1136 Engelsens, A., Hulth, S., Pihl, L., Sundbäck, K., 2008. Benthic trophic status and nutrient  
1137 fluxes in shallow-water sediments. *Estuarine, Coastal and Shelf Science* 78, 783-795.
- 1138 Engstedt, O., Stenroth, P., Larsson, P., Ljunggren, L., Elfman, M., 2010. Assessment of  
1139 natal origin of pike (*Esox lucius*) in the Baltic Sea using Sr:Ca in otoliths. *Environmental*  
1140 *Biology of Fishes* 89, 547-555.

- 1141 Engström-Öst, J., Mattila, J., 2008. Foraging, growth and habitat choice in turbid water:  
1142 an experimental study with fish larvae in the Baltic Sea. *Marine Ecology Progress Series*  
1143 359, 275-281.
- 1144 Engström-Öst, J., Immonen, E., Candolin, U., Mattila, J., 2007. The indirect effects of  
1145 eutrophication on habitat choice and survival of fish larvae in the Baltic Sea. *Marine*  
1146 *Biology* 151, 393-400.
- 1147 Eriksson, B.K., Johansson, G., Snoeijs, P., 1998. Long-term changes in the sublittoral  
1148 zonation of brown algae in the southern Bothnian Sea. *European Journal of Phycology*  
1149 33, 241-249.
- 1150 Eriksson, B.K., Sandström, A., Isæus, M., Schreiber, H., Karås, P., 2004. Effects of  
1151 boating activities on aquatic vegetation in the Stockholm archipelago, Baltic Sea.  
1152 *Estuarine, Coastal and Shelf Science* 61, 339-349.
- 1153 Eriksson, B.K., Ljunggren, L., Sandström, A., Johansson, G., Mattila, J., Rubach, A.,  
1154 Råberg, S., Snickars, M., 2009. Declines in predatory fish promote bloom-forming  
1155 macroalgae. *Ecological Applications* 19, 1975-1988.
- 1156 Eriksson B.K., Sieben, K., Eklöf, J., Ljunggren, L., Olsson, J., Casini, M., Bergström, U.,  
1157 2011. Effects of altered offshore food webs on coastal ecosystems emphasize the need  
1158 for cross-ecosystem management. *Ambio* 40, 786-797.
- 1159 Fabi, G., Spagnolo, A., Bellan-Santini, D., Charbonnel, E., Cicek, B.A., Goutayer Garcia,  
1160 J.J., Jensen, A.C., Kallianiotis, A., dos Santos, M.N., 2011. Overview on artificial reefs in  
1161 Europe. *Brazilian Journal of Oceanography* 59, 155-166.
- 1162 Florin, A.-B., Sundblad, G., Bergström, U., 2009. Characterisation of juvenile flatfish  
1163 habitats in the Baltic Sea. *Estuarine, Coastal and Shelf Science* 82, 294-300.
- 1164 Fluharty, D., 2000. Habitat protection, ecological issues, and implementation of the  
1165 Sustainable Fisheries Act. *Ecological Applications* 10, 325-337.

- 1166 Fodrie, F.J., Levin, L.A., 2008. Linking juvenile habitat utilization to population  
1167 dynamics of California halibut. *Limnology and Oceanography* 53, 799-812.
- 1168 Fodrie, F.J., Levin, L.A., Lucas, A.J., 2009. Use of population fitness to evaluate the  
1169 nursery function of juvenile habitats. *Marine Ecology Progress Series* 385, 39-49.
- 1170 Fredriksson, R., Bergström, U., Olsson, J., 2013. Riktlinjer för uppföljning av  
1171 fiskevårdsåtgärder i kustmynnande våtmarker med fokus på gädda. *Aqua Reports*  
1172 2013:7. Sveriges Lantbruksuniversitet, Öregrund, 52 pp. (In Swedish)
- 1173 Gagnon, K., Boström, C., 2016. Habitat expansion of the Harris mud crab  
1174 *Rhithropanopeus harrisii* (Gould, 1841) in the northern Baltic Sea: potential  
1175 consequences for the eelgrass food web. *BioInvasions Records* 5, 101-106.
- 1176 Gerbersdorf, S.U., Meyercordt, J., Meyer-Reil, L.A., 2005. Microphytobenthic primary  
1177 production in the Bodden estuaries, southern Baltic Sea, at two study sites differing in  
1178 trophic status. *Aquatic Microbial Ecology* 41, 181-198.
- 1179 Gibson, R.N., 1994. Impact of habitat quality and quantity on the recruitment of  
1180 juvenile flatfishes. *Netherlands Journal of Sea Research* 32, 191-206.
- 1181 Gillanders, B., Able, K., Brown, J., Eggleston, D., Sheridan, P., 2003. Evidence of  
1182 connectivity between juvenile and adult habitats for mobile marine fauna: an important  
1183 component of nurseries. *Marine Ecology Progress Series* 247, 281-295.
- 1184 Graiff, A., Liesner, D., Karsten, U., Bartsch, I., 2015. Temperature tolerance of western  
1185 Baltic Sea *Fucus vesiculosus* – growth, photosynthesis and survival. *Journal of*  
1186 *Experimental Marine Biology and Ecology* 471, 8-16.
- 1187 Graiff, A., Dankworth, M., Wahl, M., Karsten, U., Bartsch, I., 2017. Seasonal variations of  
1188 *Fucus vesiculosus* fertility under ocean acidification and warming in the western Baltic  
1189 Sea. *Botanica Marina* 60, 239-255.

- 1190 de Groot, R., Brander, L., van der Ploeg, S., Costanza, R., Bernard, F., Braat, L., Christie,  
1191 M., Crossman, N., Ghermandi, A., Hein, L., Hussain, S., Kumar, P., McVittie, A., Portela, R.,  
1192 Rodriguez, L.C., ten Brink, P., van Beukering, P., 2012. Global estimates of the value of  
1193 ecosystems and their services in monetary units. *Ecosystem Services* 1, 50-61.
- 1194 Halpern, B.S., Gaines, S.D., Warner, R.R., 2005. Habitat size, recruitment, and longevity  
1195 as factors limiting populations size in stage-structured species. *American Naturalist* 165,  
1196 82-94.
- 1197 Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno,  
1198 J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P.,  
1199 Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A global map of human  
1200 impact on marine ecosystems. *Science* 319, 948-952.
- 1201 Hansen, J.P., Snickars, M., 2014. Applying macrophyte community indicators to assess  
1202 anthropogenic pressures on shallow soft bottoms. *Hydrobiologia* 738, 171-189.
- 1203 Härmä, M., Lappalainen, A., Urho, L., 2008. Reproduction areas of roach (*Rutilus*  
1204 *rutilus*) in the northern Baltic Sea: potential effects of climate change. *Canadian Journal*  
1205 *of Fisheries and Aquatic Sciences* 65, 2678-2688.
- 1206 Havenhand, J.N., 2012. How will ocean acidification affect Baltic Sea ecosystems? An  
1207 assessment of plausible impacts on key functional groups. *Ambio* 41, 637-644.
- 1208 Heikinheimo, O., Pekcan-Hekim, Z., Raitaniemi, J., 2014. Spawning stock–recruitment  
1209 relationship in pikeperch *Sander lucioperca* (L.) in the Baltic Sea, with temperature as an  
1210 environmental effect. *Fisheries Research* 155, 1-9.
- 1211 HELCOM, 2007. HELCOM Baltic Sea Action Plan. Helsinki Commission.
- 1212 HELCOM, 2010. Towards a tool for quantifying the anthropogenic pressures and  
1213 potential impacts in the Baltic Sea marine environment. A background document on the  
1214 method, data preparation and testing of the Baltic Sea Pressure and Impact Indices.

- 1215 Baltic Sea Environment Proceedings 125. Available at: <http://www.helcom.fi/helcom->  
1216 [at-work/publications/baltic-sea-environment-proceedings/](http://www.helcom.fi/helcom-at-work/publications/baltic-sea-environment-proceedings/)
- 1217 HELCOM, 2012. Indicator based assessment of coastal fish community status in the  
1218 Baltic Sea 2005-2009. Baltic Sea Environment Proceeding No. 131.
- 1219 HELCOM, 2013. Climate change in the Baltic Sea Area: HELCOM thematic assessment  
1220 in 2013. Baltic Sea Environment Proceeding No. 137.
- 1221 HELCOM, 2017, First version of the 'State of the Baltic Sea' report – June 2017 – to be  
1222 updated in 2018. Available at: <http://stateofthebalticsea.helcom.fi>
- 1223 Helle, I., Jolma, A., Venesjärvi, R., 2016. Species and habitats in danger: estimating the  
1224 relative risk posed by oil spills in the northern Baltic Sea. *Ecosphere* 7, e01344
- 1225 Hemminga, M.A., Duarte, C.M., 2000. *Seagrass Ecology*. Cambridge University Press,  
1226 Cambridge, 298 pp.
- 1227 Hinrichsen, H.H., von Dewitz, B., Lehmann, A., Bergström, U., Hüsey, K. (2017). Spatio-  
1228 temporal dynamics of cod nursery areas in the Baltic Sea. *Progress in Oceanography*  
1229 155, 28-40.
- 1230 Holmlund, C.M., Hammer, M., 1999. Ecosystem services generated by fish populations.  
1231 *Ecological Economics* 29, 253-268.
- 1232 Hopkins, C.C.E., 2003. The dangers of bottom trawling in the Baltic Sea. A report for  
1233 Coalition Clean Baltic. Aquamarine advisers.
- 1234 Hopkins, T.S., Bailly, D., Støttrup, J.G., 2011. A Systems Approach Framework for  
1235 Coastal Zones. *Ecology and Society* 16, 25.
- 1236 Idestam-Almquist, J., 2000. Dynamics of submersed aquatic vegetation on shallow soft  
1237 bottoms in the Baltic Sea. *Journal of Vegetation Science* 11, 425-432.
- 1238 Iles, T.C., Beverton, R.J.H., 2000. The concentration hypothesis: the statistical  
1239 evidence. *ICES Journal of Marine Science* 57, 216-227.

- 1240 Islam, M.S., Tanaka, M., 2004. Impacts of pollution on coastal and marine ecosystems  
1241 including coastal and marine fisheries and approach for management: a review and  
1242 synthesis. *Marine Pollution Bulletin* 48, 624-649.
- 1243 Ivarsson, M., Magnussen, K., Heiskanen, A.-S., Navrud, S., Viitasalo, M., 2017.  
1244 Ecosystem services in MSP: Ecosystem services approach as a common Nordic  
1245 understanding for MSP. *TemaNord* 2017:536. Nordic Council of Ministers.  
1246 <http://dx.doi.org/10.6027/TN2017-56>
- 1247 Jackson, E.L., Rowden, A.A., Attrill, M.J., Bossey, S.J., Jones, M.B., 2001. The importance  
1248 of seagrass beds as a habitat for fishery species. *Oceanography and Marine Biology: an*  
1249 *Annual Review* 39, 269-304.
- 1250 Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J.,  
1251 Bradbury, R., Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B.,  
1252 Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., Warner, R., 2001.  
1253 Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293, 629-  
1254 638.
- 1255 Järv, L., Kotta, J., Kotta, I., Raid, T., 2011. Linking the structure of benthic invertebrate  
1256 communities and the diet of native and invasive fish species in a brackish water  
1257 ecosystem. *Annales Zoologici Fennici* 49, 129-141.
- 1258 Jokinen, H., Wennhage, H., Lappalainen, A., Ådjers, K., Rask, M., Norkko, A., 2015.  
1259 Decline of flounder (*Platichthys flesus* (L.)) at the margin of the species' distribution  
1260 range. *Journal of Sea Research* 105, 1-9.
- 1261 Jokinen, H., Wennhage, H., Ollus, V., Aro, E., Norkko, A., 2016. Juvenile flatfish in the  
1262 northern Baltic Sea - long-term decline and potential links to habitat characteristics.  
1263 *Journal of Sea Research* 107, 67-75.

- 1264 Jormalainen, V., Gagnon, K., Sjöroos, J., Rothäusler, E., 2016. The invasive mud crab  
1265 enforces a major shift in a rocky littoral invertebrate community of the Baltic Sea.  
1266 *Biological Invasions* 18, 1409-1419.
- 1267 Kallasvuoto, M., Salonen, M., Lappalainen, A., 2009. Does the zooplankton prey  
1268 availability limit the larval habitats of pike in the Baltic Sea? *Estuarine, Coastal and Shelf*  
1269 *Science* 86, 148-156.
- 1270 Kallasvuoto, M., Vanhatalo, J., Veneranta, L., 2017. Modeling the spatial distribution of  
1271 larval fish abundance provides essential information for management. *Canadian Journal*  
1272 *of Fisheries and Aquatic Sciences* 74, 636-649.
- 1273 Kappel, C.V., 2005. Losing pieces of the puzzle: threats to marine, estuarine, and  
1274 diadromous species. *Frontiers in Ecology and the Environment* 3, 275-282.
- 1275 Karås, P., 1996. Basic abiotic conditions for production of perch (*Perca fluviatilis* L.)  
1276 young-of-the-year in the Gulf of Bothnia. *Annales Zoologici Fennici* 33, 371-381.
- 1277 Karås, P., Hudd, R., 1993. Reproduction areas of fish in the northern Quark (Gulf of  
1278 Bothnia). *Aqua Fennica* 23, 39-49.
- 1279 Karlson, A.M.L., Almqvist, G., Skora, K.E., Appelberg, M., 2007. Indications of  
1280 competition between non-indigenous round goby and native flounder in the Baltic Sea.  
1281 *ICES Journal of Marine Science* 64, 479-486.
- 1282 Karlson, K., Rosenberg, R., Bonsdorff, E., 2002. Temporal and spatial large-scale effects  
1283 of eutrophication and oxygen deficiency on benthic fauna in Scandinavian and Baltic  
1284 waters – a review. *Oceanography Marine Biology: an Annual Review* 40, 427-489.
- 1285 Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çinar, M.E., Öztürk, B.,  
1286 Grabowski, M., Golani, D., Cardoso, A.C., 2014. Impacts of invasive alien marine species  
1287 on ecosystem services and biodiversity: a pan-European review. *Aquatic Invasions* 9,  
1288 391-423.



- 1289 Kautsky, N., Kautsky, H., Kautsky, U., Waern, M., 1986. Decreased depth penetration of  
1290 *Fucus vesiculosus* (L.) since the 1940's indicates eutrophication of the Baltic Sea. Marine  
1291 Ecology Progress Series 28, 1-8.
- 1292 Kenny, A.J., Skjoldal, H.R., Engelhard, G.H., Kershaw, P.J., Reid, J.B., 2009. An integrated  
1293 approach for assessing the relative significance of human pressures and environmental  
1294 forcing on the status of Large Marine Ecosystems. Progress in Oceanography 81, 132-  
1295 148.
- 1296 Kneib, K.T., Wagner, S.L., 1994. Nekton use of vegetated marsh habitats at different  
1297 stages of tidal inundation. Marine Ecology Progress Series 106, 227-238.
- 1298 Kornis, M.S., Mercado-Silva, N., Vander Zanden, M.J., 2012. Twenty years of invasion: a  
1299 review of round goby *Neogobius melanostomus* biology, spread and ecological  
1300 implications. Journal of Fish Biology 80, 235-285.
- 1301 Korpinen, S., Honkanen, T., Vesakoski, O., Hemmi, A., Koivikko, R., Lojonen, J.,  
1302 Jormalainen, V., 2007. Macroalgal communities face the challenge of changing biotic  
1303 interactions: review with focus on the Baltic Sea. Ambio 36, 203-211.
- 1304 Korpinen, S., Meski, L., Andersen, J.H., Laamanen, M., 2012. Human pressures and their  
1305 potential impact on the Baltic Sea ecosystem. Ecological Indicators 15, 105-114.
- 1306 Kotta, J., Nurkse, K., Puntila, R., Ojaveer, H., 2016. Shipping and natural environmental  
1307 conditions determine the distribution of the invasive non-indigenous round goby  
1308 *Neogobius melanostomus* in a regional sea. Estuarine, Coastal and Shelf Science 169, 15-  
1309 24.
- 1310 Kraufvelin, P., 2007. Responses to nutrient enrichment, wave action and disturbance  
1311 in rocky shore communities. Aquatic Botany 87, 262-274.
- 1312 Kraufvelin, P., Díaz, E.R., 2015. Sediment macrofauna communities at a small mussel  
1313 farm in the northern Baltic proper. Boreal Environment Research 20, 378-390.

- 1314 Kraufvelin, P., Ruuskanen, A.T., Nappu, N., Kiirikki, M., 2007. Winter colonisation and  
1315 succession of filamentous algae on artificial substrates and possible relationships to  
1316 *Fucus vesiculosus* settlement in early summer. Estuarine, Coastal and Shelf Science 72,  
1317 665-674.
- 1318 Kraufvelin, P., Lindholm, A., Pedersen, M.F., Kirkerud, L.A., Bonsdorff, E., 2010.  
1319 Biomass, diversity and production of rocky shore macroalgae at two nutrient  
1320 enrichment and wave action levels. Marine Biology 157, 29-47.
- 1321 Kraufvelin, P., Ruuskanen, A.T., Bäck, S., Russell, G., 2012. Increased seawater  
1322 temperature and light during early springs accelerate receptacle growth of *Fucus*  
1323 *vesiculosus* in the northern Baltic proper. Marine Biology 159, 1795-1807.
- 1324 Kraufvelin, P., Pekcan-Hekim, Z., Bergström, U., Florin, A.-B., Lehtikainen, A., Mattila, J.,  
1325 Olsson, J., 2016. Essential fish habitats (EFH): Conclusions from a workshop on the  
1326 importance, mapping, monitoring, threats and conservation of coastal EFH in the Baltic  
1327 Sea. TemaNord 2016, 539.
- 1328 Lappalainen, A., Westerborn, M., Vesala, S., 2004. Blue mussels (*Mytilus edulis*) in the  
1329 diet of roach (*Rutilus rutilus*) in outer archipelago areas of the western Gulf of Finland,  
1330 Baltic Sea. Hydrobiologia 514, 87-92.
- 1331 Lappalainen, A., Westerborn, M., Heikinheimo, O., 2005. Roach (*Rutilus rutilus*) as an  
1332 important predator on blue mussel (*Mytilus edulis*) populations in a brackish water  
1333 environment, the northern Baltic Sea. Marine Biology 147, 323-330.
- 1334 Lappalainen, A., Härma, M., Kuningas, S., Urho, L., 2008. Reproduction of pike (*Esox*  
1335 *lucius*) in reed belt shores of the SW coast of Finland, Baltic Sea: a new survey approach.  
1336 Boreal Environment Research 13, 370-380.
- 1337 Laurila-Pant, M., Lehtikainen, A., Uusitalo, L., Venesjärvi, R., 2015. How to value  
1338 biodiversity in environmental management? Ecological Indicators 55, 1-11.

- 1339 Leathwick, J., Moilanen, A., Francis, M., Elith, J., Taylor, P., Julian, K., Hastie, T., Duffy, C.,  
1340 2008. Novel methods for the design and evaluation of marine protected areas in offshore  
1341 waters. *Conservation Letters* 1, 91-102.
- 1342 Leppäkoski, E., Bonsdorff, E., 1989. Ecosystem variability and gradients. Examples  
1343 from the Baltic Sea as a background for hazard assessment. In: Landner, L. (Ed.),  
1344 *Chemicals in the Aquatic Environment*, Springer Berlin Heidelberg, pp. 6-58.
- 1345 Leppäkoski, E., Gollasch, S., Gruszka, P., Ojaveer, H., Olenin, S., Panov, V., 2002. The  
1346 Baltic a sea of invaders. *Canadian Journal of Fisheries and Aquatic Sciences* 59, 1175-  
1347 1188.
- 1348 Levin, P.S., Stunz, G.W., 2005. Habitat triage for exploited fishes: Can we identify  
1349 essential "Essential Fish Habitat?" *Estuarine, Coastal and Shelf Science* 64, 70-78.
- 1350 Liversage, K., Nurkse, K., Kotta, J., Järv, L., 2017. Environmental heterogeneity  
1351 associated with European perch (*Perca fluviatilis*) predation on invasive round goby  
1352 (*Neogobius melanostomus*). *Marine Environmental Research* 132, 132-139.
- 1353 Ljunggren, L., 2002. Growth response of pikeperch larvae in relation to body size and  
1354 zooplankton abundance. *Journal of Fish Biology* 60, 405-414.
- 1355 Ljunggren, L., Sandström A., Bergström U., Mattila J., Lappalainen A., Johansson G.,  
1356 Sundblad G., Casini M., Kaljuste O., Eriksson, B.K., 2010. Recruitment failure of coastal  
1357 predatory fish in the Baltic Sea is related to an offshore system shift. *ICES Journal of*  
1358 *Marine Science* 67, 1587-1595.
- 1359 Lokko, K., Kotta, J., Orav-Kotta, H., Nurkse, K., Pärnoja, M., 2017. Introduction of a  
1360 functionally novel consumer to a low diversity system: effects of the mud crab  
1361 *Rhithropanopeus harrisii* on meiobenthos. *Estuarine, Coastal and Shelf Science* (in press).
- 1362 Long, R.D., Charles, A., Stephenson, R.L., 2015. Key principles of marine ecosystem-  
1363 based management. *Marine Policy* 57, 53-60.

Lotze, H.K., 2004. Repetitive history of resource depletion and mismanagement: the need for a shift in perspective. *Marine Ecology Progress Series* 274, 269-303.

Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H, Cooke, R.G., Kay, M.C., Kidwell, S.M., Kirby, M.X., Peterson, C.H., Jackson, J.B.C., 2006. Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science* 312, 1806-1809.

Lundberg, C., 2005. Eutrophication in the Baltic Sea: From area-specific biological effects to interdisciplinary consequences. PhD-thesis, Åbo Akademi University, 166 pp.

MacKenzie, B.R., Hinrichsen, H.H., Plikshs, M., Wieland, K., Zezera, A.S., 2000. Quantifying environmental heterogeneity: estimating the size of habitat for successful cod egg development in the Baltic Sea. *Marine Ecology Progress Series* 193, 143–156.

MacKenzie, B.R., Gislason, H., Möllmann, C., Köster, F.W., 2007. Impact of 21st century climate change on the Baltic Sea fish community and fisheries. *Global Change Biology*, 13, 1348-1367.

Macura, B., Lönnstedt, O.M., Byström, P., Airolidi, L., Eriksson, B.K., Rudstam, L., Støttrup, J., 2016. What is the impact on fish recruitment of anthropogenic physical and structural habitat change in shallow nearshore areas in temperate systems? A systematic review protocol. *Environmental Evidence* 5, 10.

Mangel, M., Levin, P., Patil, A., 2006. Using life history and persistence criteria to prioritize habitats for management and conservation. *Ecological Applications* 16, 797-806.

McCormick, S.D., Hansen, L.P., Quinn, T.P., Saunders, R.L., 1998. Movement, migration, and smolting of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* 55, 77-92.

McLusky, D.S., Elliott, M., 2004. *The Estuarine Ecosystem: Ecology, Threats and Management*. New York: Oxford University Press. 224.

- 1389 Meriste, M., Kirsimäe, K., 2015. Development of the reed bed in Matsalu wetland,  
1390 Estonia: responses to neotectonic land uplift, sea level changes and human influences.  
1391 Estonian Journal of Earth Sciences 64, 159.
- 1392 Minns, C.K., Randall, R.G., Moore, J.E., Cairns, V.W., 1996. A model simulating the  
1393 impact of habitat supply limits on northern pike, *Esox lucius*, in Hamilton Harbour, Lake  
1394 Ontario. Canadian Journal of Fisheries and Aquatic Sciences 53, 20-34.
- 1395 Möllmann, C., Lindegren, M., Blenckner, T., Bergström, L., Casini, M., Diekmann, R.,  
1396 Flinkman, J., Müller-Karulis, B., Neuenfeldt, S., Schmidt, J.O., Tomczak, M., Voss, R.,  
1397 Gårdmark, A., 2014. Implementing ecosystem-based fisheries management: from single-  
1398 species to integrated ecosystem assessment and advice for Baltic Sea fish stocks. ICES  
1399 Journal of Marine Science 71, 1187-1197.
- 1400 Moore, C., Drazen, J.C., Radford, B.T., Kelley, C., Newman, S.J., 2016. Improving  
1401 essential fish habitat designation to support sustainable ecosystem-based fisheries  
1402 management. Marine Policy 69, 32-41.
- 1403 Mumby, P.J., Edwards, A.J., Arias-González, J.E., Lindeman, K.C., Blackwell, P.G., Gall, A.,  
1404 Gorczynska, M.I., Harborne, A.R., Pescod, C.L., Renken, H., Wabnitz, C.C.C., Llewellyn, G.,  
1405 2004. Mangroves enhance the biomass of coral reef fish communities in the Caribbean.  
1406 Nature 427, 533-536.
- 1407 Newell, R.C., Seiderer, L.J., Hitchcock, D.R., 1998. The impact of dredging works in  
1408 coastal waters: a review of the sensitivity to disturbance and subsequent recovery of  
1409 biological resources on the sea bed. Oceanography and Marine Biology: an Annual  
1410 Review 36, 127-178.
- 1411 Nilsson, J., Engstedt, O., Larsson, P., 2014. Wetlands for northern pike (*Esox lucius* L.)  
1412 recruitment in the Baltic Sea. Hydrobiologia 721, 145-154.

- 1413 Nurkse, K., Kotta, J., Orav-Kotta, H., Pärnoja, M., Kuprijanov, I., 2015. Laboratory  
1414 analysis of the habitat occupancy of the crab *Rhithropanopeus harrisii* (Gould) in an  
1415 invaded ecosystem: the north-eastern Baltic Sea. *Estuarine, Coastal and Shelf Science*,  
1416 154, 152-157.
- 1417 Nurkse, K., Kotta, J., Orav-Kotta, H., Ojaveer, H., 2016. A successful non-native  
1418 predator, round goby, in the Baltic Sea: generalist feeding strategy, diverse diet and high  
1419 prey consumption. *Hydrobiologia* 777, 271-281.
- 1420 Oesterwind, D., Bock, C., Förster, A., Gabel, M., Henseler, C., Kotterba, P., Menge, M.,  
1421 Myts, D., Winkler, H.M., 2017. Predator and prey: the role of the round goby *Neogobius*  
1422 *melanostomus* in the western Baltic. *Marine Biology Research* 13, 188-197.
- 1423 Ojaveer, H., Kotta, J., 2015. Ecosystem impacts of the widespread non-indigenous  
1424 species in the Baltic Sea: literature survey evidences major limitations in knowledge.  
1425 *Hydrobiologia* 750, 171-185.
- 1426 Ojaveer, H., Galil, B.S., Lehtiniemi, M., Christoffersen, M., Clink, S., Florin, A.-B., Gruszka,  
1427 P., Puntila, R., Behrens, J.W., 2015. Twenty five years of invasion: management of the  
1428 round goby *Neogobius melanostomus* in the Baltic Sea. *Management of Biological*  
1429 *Invasions* 6, 329-339.
- 1430 Olsson, J., Bergström, L., Gårdmark, A., 2012. Abiotic drivers of coastal fish community  
1431 change during four decades in the Baltic Sea. *ICES Journal of Marine Science* 69, 961-  
1432 970.
- 1433 Olsson, J., Bergström, L., Gårdmark, A., 2013. Top-down regulation, climate and multi-  
1434 decadal changes in coastal zoobenthos communities in two Baltic Sea areas. *PLoS One*  
1435 8:e64767.
- 1436 Olsson, J., Tomczak, M.T., Ojaveer, H., Gårdmark, A., Pöllumäe, A., Müller-Karulis, B.,  
1437 Ustups, D., Dinesen, G.E., Peltonen, H., Putnis, I., Szymanek, L., Simm, M., Heikinheimo, O.,

- 1438 Gasyukov, P., Axe, P., Bergström, L., 2015. Temporal development of coastal ecosystems  
1439 in the Baltic Sea over the past two decades. ICES Journal of Marine Science 72, 2539-  
1440 2548.
- 1441 Orio, A., Bergström, U., Casini, M., Erlandsson, M., Eschbaum, R., Hüsey, K., Lehmann,  
1442 A., Ložys L., Ustups, D., Florin, A-B., 2017. Characterizing and predicting the distribution  
1443 of Baltic Sea flounder (*Platichthys flesus*) during the spawning season. Journal of Sea  
1444 Research 126, 46-55.
- 1445 Orth, R.J., Carruthers, T.J., Dennison, W.C., Duarte, C.M., Fourqurean, J.W., Heck, K.L.,  
1446 Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Olyarnik, S., Short, F.T., Waycott, M.,  
1447 Williams, S.L., 2006. A global crisis for seagrass ecosystems. Bioscience 56, 987-996.
- 1448 Österblom, H., Hansson, S., Larsson, U., Hjerne, O., Wulff, F., Elmgren, R., Folke, C.,  
1449 2007. Human-induced trophic cascades and ecological regime shifts in the Baltic Sea.  
1450 Ecosystems 10, 877-889.
- 1451 Östman, Ö., Eklöf, J., Eriksson, B.K., Olsson, J., Moksnes, P.-O., Bergström, U., 2016. Top-  
1452 down control as important as nutrient enrichment for eutrophication effects in North  
1453 Atlantic coastal ecosystems. Journal of Applied Ecology 53, 1138-1147.
- 1454 Östman, Ö., Lingman, A., Bergström, L., Olsson, J., 2017. Temporal development and  
1455 spatial scale of coastal fish indicators in reference ecosystems: hydroclimate and  
1456 anthropogenic drivers. Journal of Applied Ecology 54, 557-566.
- 1457 Paerl, H.W., Otten, T.G., 2013. Harmful cyanobacterial blooms: causes, consequences,  
1458 and controls. Microbial Ecology 65, 995-1010.
- 1459 Persson, L., Diehl, S., Johansson, L., Andersson, G., Hamrin, S.F., 1991. Shifts in fish  
1460 communities along the productivity gradient of temperate lakes – patterns and the  
1461 importance of size-structured interactions. Journal of Fish Biology 38, 281-293.

- 1462 Pihl, L., Wennhage, H., 2002. Structure and diversity of fish assemblages on rocky and  
1463 soft bottom shores on the Swedish west coast. *Journal of Fish Biology* 61, 148-166.
- 1464 Pihl, L., Wennhage, H., Nilsson S., 1994. Fish assemblage structure in relation to  
1465 macrophytes and filamentous epiphytes in shallow non-tidal rocky- and soft-bottom  
1466 habitats. *Environmental Biology of Fishes* 39, 271-288.
- 1467 Pikitch, E.K., Santora, C., Babcock, E.A., Bakun, A., Bonfil, R., Conover, D.O., Dayton, P.K.,  
1468 Doukakis, P., Fluharty, D., Heneman, B., Houde, E.D., Link, J., Livingston, P.A., Mangel, M.,  
1469 McAllister, M.K., Pope, J., Sainsbury, K.J., 2004. Ecosystem-based fishery management.  
1470 *Science* 305, 346-347.
- 1471 Pikitch, E.K., Rountos, K.J., Essington, T.E., Santora, C., Pauly, D., Watson, R., Sumaila,  
1472 U.R., Boersma, P.D., Boyd, I.L., Conover, D.O., Cury, P., Heppell, S.S., Houdes, E.D., Mangel,  
1473 M., Plagányi, E., Sainsbury, K., Steneck, R.S., Geers, T.M., Gownaris, Munch, S.B., 2014. The  
1474 global contribution of forage fish to marine fisheries and ecosystems. *Fish and Fisheries*  
1475 15, 43-64.
- 1476 Pitkänen, H., Peuraniemi, M., Westerborn, M., Kilpi, M., von Numers, M., 2013. Long-  
1477 term changes in distribution and frequency of aquatic vascular plants and charophytes  
1478 in an estuary in the Baltic Sea. *Annales Botanici Fennici* 50, 1-54.
- 1479 Pittman, S.J., Brown, K.A., 2011. Multi-scale approach for predicting fish species  
1480 distributions across coral reef seascapes. *PLoS One* 6, e20583.
- 1481 Plinski, M., Florczyk, I., 1984. Changes in the phytobenthos resulting from the  
1482 eutrophication of the Puck Bay. *Limnologia* 15, 325-327.
- 1483 Pommer, C.D., Olesen, M., Hansen, J.L., 2016. Impact and distribution of bottom trawl  
1484 fishing on mud-bottom communities in the Kattegat. *Marine Ecology Progress Series*  
1485 548, 47-60.



- 1486 Powers, S.P., Peterson, C.H., Christian, R.R., Sullivan, E., Powers, M.J., Bishop, M.J.,  
1487 Buzzelli, C.P., 2005. Effects of eutrophication on bottom habitat and prey resources of  
1488 demersal fishes. *Marine Ecology Progress Series* 302, 233-243.
- 1489 Pulkkinen, H., Mäntyniemi, S., Kuikka, S., Levontin, P., 2011. More knowledge with the  
1490 same amount of data: advantage of accounting for parameter correlations in hierarchical  
1491 meta-analyses. *Marine Ecology Progress Series* 443, 29-37.
- 1492 Råberg, S., Jönsson B.R., Björn, A., Granéli, E., Kautsky, L., 2005. Effects of *Pilayella*  
1493 *littoralis* on *Fucus vesiculosus* recruitment. Implications for community composition.  
1494 *Marine Ecology Progress Series* 289, 131-139.
- 1495 Rahikainen, M., Helle, I., Haapasaari, P., Oinonen, S., Kuikka, S., Vanhatalo, J.,  
1496 Mäntyniemi, S., Hoviniemi, K.M., 2014. Toward integrative management advice of water  
1497 quality, oil spills, and fishery in the Gulf of Finland: A Bayesian approach. *Ambio* 43, 115-  
1498 123.
- 1499 Rajasilta, M., Eklund, J., Kääriä, J., Ranta-Aho, K., 1989. The deposition and mortality of  
1500 the eggs of the Baltic herring *Clupea harengus membras* L., on different substrates in the  
1501 south-west archipelago of Finland. *Journal of Fish Biology* 34, 417-427.
- 1502 Rakauskas, V., Pūtys, Ž., Dainys, J., Lesutienė, J., Ložys, L., Arbačiauskas, K., 2013.  
1503 Increasing population of the invader round goby, *Neogobius melanostomus*  
1504 (Actinopterygii: Perciformes: Gobiidae), and its trophic role in the Curonian Lagoon, SE  
1505 Baltic Sea. *Acta Ichthyologica et Piscatoria* 43, 95-108.
- 1506 Rice, J.C., 2005. Understanding fish habitat ecology to achieve conservation. *Journal of*  
1507 *Fish Biology* 67, 1-22.
- 1508 Rijnsdorp, A.D., Van Beek, F.A., Flatman, S., Millner, R.M., Riley, J.D., Giret, M., De  
1509 Clerck, R., 1992. Recruitment of sole stocks, *Solea solea* (L.), in the Northeast Atlantic.  
1510 *Netherlands Journal of Sea Research* 29, 173-192.

- 1511 Rinne, H., Salovius-Laurén, S., Mattila, J., 2011. The occurrence and depth penetration  
1512 of macroalgae along environmental gradients in the northern Baltic Sea. *Estuarine,*  
1513 *Coastal and Shelf Science* 94, 182-191.
- 1514 Rohtla, M., 2015. Otolith sclerochronological studies on migrations, spawning habitat  
1515 preferences and age of freshwater fishes inhabiting the Baltic Sea. Doctoral dissertation,  
1516 University of Tartu, Estonia.
- 1517 Rohtla, M., Vetemaa, M., Urtson, K., Soesoo, A., 2012. Early life migration patterns of  
1518 Baltic Sea pike *Esox lucius*. *Journal of Fish Biology* 80, 886-893.
- 1519 Rohtla, M., Vetemaa, M., Taal, I., Svirgsden, R., Urtson, K., Saks, L., Verliin, A., Kesler, M.,  
1520 Saat, T., 2014. Life history of anadromous burbot (*Lota lota*, Linneaus) in the brackish  
1521 Baltic Sea inferred from otolith microchemistry. *Ecology of Freshwater Fish* 23, 141-  
1522 148.
- 1523 Rohtla, M., Svirgsden, R., Taal, I., Saks, L., Eschbaum, R., Vetemaa, M., 2015. Life history  
1524 characteristics of ide *Leuciscus idus* in the Eastern Baltic Sea. *Fisheries Management and*  
1525 *Ecology* 22, 239-248.
- 1526 Rohtla, M., Svirgsden, R., Verliin, A., Rumvolt, K., Matetski, L., Hommik, K., Saks, L.,  
1527 Vetemaa, M., 2017. Developing novel means for unravelling population structure,  
1528 provenance and migration patterns of European whitefish *Coregonus lavaretus* s.l. in the  
1529 Baltic Sea. *Fisheries Research* 187, 47-57.
- 1530 Rönnbäck, P., Kautsky, N., Pihl, L., Troell, M., Söderqvist, T., Wennhage, H., 2007.  
1531 Ecosystem goods and services from Swedish coastal habitats: identification, valuation,  
1532 and implications of ecosystem shifts. *Ambio* 36, 534-544
- 1533 Rose, K.A., 2000. Why are quantitative relationships between environmental quality  
1534 and fish populations so elusive? *Ecological Applications* 10, 367-385.

- 1535 Rosenberg, A., Bigford, T.E., Leathery, S., Hill, R.L., Bickers, K., 2000. Ecosystem  
1536 approaches to fishery management through essential fish habitat. *Bulletin of Marine*  
1537 *Science* 66, 535-542.
- 1538 Rosqvist, K., Mattila, J., Sandström, A., Snickars, M., Westerborn, M., 2010. Regime  
1539 shifts in vegetation composition of Baltic Sea coastal lagoons. *Aquatic Botany* 93, 39-46.
- 1540 Sandström, A., Karås, P., 2002. Effects of eutrophication on young-of-the-year  
1541 freshwater fish communities in coastal areas of the Baltic. *Environmental Biology of*  
1542 *Fishes* 63, 89-101.
- 1543 Sandström, A., Eriksson, B.K., Karås, P., Isæus, M., Schreiber, H., 2005. Boating and  
1544 navigation activities influence the recruitment of fish in a Baltic Sea archipelago area.  
1545 *Ambio* 34, 125-130.
- 1546 Schmitt, R.J., Holbrook, S.J., 2000. Habitat-limited recruitment of coral reef damselfish.  
1547 *Ecology* 81, 3479-3494.
- 1548 Seaman, W., 2007. Artificial habitats and the restoration of degraded marine  
1549 ecosystems and fisheries. *Hydrobiologia* 580, 143-155.
- 1550 Seitz, R.D., Wennhage, H., Bergström, U., Lipcius, R.N., Ysebaert, T., 2014. Ecological  
1551 value of coastal habitats for commercially and ecologically important species. *ICES*  
1552 *Journal of Marine Science* 71, 648-665.
- 1553 Sheaves, M., Baker, R., Nagelkerken, I., Connolly, R.M., 2015. True value of estuarine  
1554 and coastal nurseries for fish: Incorporating complexity and dynamics. *Estuaries and*  
1555 *Coasts* 38, 401-414.
- 1556 Šiaulys, A., Daunys, D., Bučas, M. and Bacevičius, E., 2012. Mapping an ecosystem  
1557 service: A quantitative approach to derive fish feeding ground maps. *Oceanologia* 54,  
1558 491-505.

- 1559 Sissenwine, M., Symes, D., 2007. Reflections on the Common Fisheries Policy. Report  
1560 to the general directorate for fisheries and maritime affairs of the European commission,  
1561 pp. 1–75.
- 1562 Snickars, M., Sandström, A., Lappalainen, A., Mattila, J., Rosqvist, K., Urho, L., 2009.  
1563 Fish assemblages in coastal lagoons in land-uplift succession: the relative importance of  
1564 local and regional environmental gradients. *Estuarine, Coastal and Shelf Science* 81, 247-  
1565 256.
- 1566 Snickars, M., Sundblad, G., Sandström, A., Ljunggren, L., Bergström, U., Johansson, G.,  
1567 Mattila, J., 2010. Habitat selectivity of substrate-spawning fish: modelling requirements  
1568 for the Eurasian perch *Perca fluviatilis*. *Marine Ecology Progress Series* 398, 235-243.
- 1569 Snickars, M., Weigel, B., Bonsdorff, E., 2015. Impact of eutrophication and climate  
1570 change on fish and zoobenthos in coastal waters of the Baltic Sea. *Marine Biology* 162,  
1571 141-151.
- 1572 Sparholt, H., 1994. Fish species interactions in the Baltic Sea. *Dana* 10, 131-162.
- 1573 Stål, J., Pihl, L., Wennhage, H., 2007. Food utilisation by coastal fish assemblages in  
1574 rocky and soft bottoms on the Swedish west coast: Inference for identification of  
1575 essential fish habitats. *Estuarine, Coastal and Shelf Science* 71, 593-607.
- 1576 Stelzenmüller, V., Lee, J., Garnacho, E., Rogers, S.I., 2010. Assessment of a Bayesian  
1577 Belief Network–GIS framework as a practical tool to support marine planning. *Marine*  
1578 *Pollution Bulletin* 60, 1743-1754.
- 1579 Støttrup, J.G., Stenberg, C., Dahl, K., Kristensen, L.D., Richardson, K., 2014. Restoration  
1580 of a temperate reef: Effects on the fish community. *Open Journal of Ecology* 4:1045-  
1581 1059.

- 1582 Støttrup, J.G., Dinesen, G.E., Janssen, H., Gillgren, C., Schernewski, G., 2016. Re-visiting  
1583 ICM theory and practice: Lessons learned from the Baltic Sea Region. *Ocean and Coastal*  
1584 *Management* 139, 64-76.
- 1585 Sundblad, G., Bergström, U., 2014. Shoreline development and degradation of coastal  
1586 fish reproduction habitats. *Ambio* 43, 1020-1028.
- 1587 Sundblad, G., Bergström, U., Sandström, A., 2011. Ecological coherence of marine  
1588 protected area networks: a spatial assessment using species distribution models. *Journal*  
1589 *of Applied Ecology* 48, 112-120.
- 1590 Sundblad, G., Bergström, U., Sandström, A., Eklöv, P., 2014. Nursery habitat availability  
1591 limits adult stock sizes of predatory coastal fish. *ICES Journal of Marine Science* 71, 672-  
1592 680.
- 1593 Thompson, R.C., Crowe, T.P., Hawkins, S.J., 2002. Rocky intertidal communities: past  
1594 environmental changes, present status and predictions for the next 25 years.  
1595 *Environmental Conservation* 29, 168-191.
- 1596 Thomsen, J., Gutowska, M.A., Saphorster, J., Heinemann, A., Trubenbach, K., Fietzke, J.,  
1597 Hiebenthal, C., Eisenhauer, A., Körtzinger, A., Wahl, M., Melzner, F., 2010. Calcifying  
1598 invertebrates succeed in a naturally CO<sub>2</sub>-rich coastal habitat but are threatened by high  
1599 levels of future acidification. *Biogeosciences* 7, 3879-3891.
- 1600 Thrush, S.F., Dayton, P.K., 2010. What can ecology contribute to ecosystem-based  
1601 management? *Annual Review of Marine Science* 2, 419-441.
- 1602 Torn, K., Herkül, K., Martin, G., Oganjan, K. (2017). Assessment of quality of three  
1603 marine benthic habitat types in northern Baltic Sea. *Ecological Indicators*, 73, 772-783.
- 1604 Turner, S.J., Thrush, S.F., Hewitt, J.E., Cummings, V.J., Funnell, G., 1999. Fishing impacts  
1605 and the degradation or loss of habitat structure. *Fisheries Management and Ecology* 6,  
1606 401-420.

- 1607 Ustups, D., Müller-Karulis, B., Bergström, U., Makarchouk, A., Sics, I., 2013. The  
1608 influence of environmental conditions on early life stages of flounder (*Platichthys flesus*)  
1609 in the central Baltic Sea. *Journal of Sea Research* 75, 77-84.
- 1610 Ustups, D., Bergström, U., Florin, A.-B., Kruze, E., Zilniece, D., Elferts, D., Knospina, E,  
1611 Uzars, D. (2016). Diet overlap between juvenile flatfish and the invasive round goby in  
1612 the central Baltic Sea. *Journal of Sea Research*, 107, 121-129.
- 1613 Uusitalo, L., Kuikka, S., Kauppila, P., Söderkultalahti, P., Bäck, S., 2012. Assessing the  
1614 roles of environmental factors in coastal fish production in the northern Baltic Sea: A  
1615 Bayesian network application. *Integrated Environmental Assessment and Management*  
1616 8, 445-455.
- 1617 Uusitalo, L., Korpinen, S., Andersen, J.H., Niiranen, S., Valanko, S., Heiskanen, A.S.,  
1618 Dickey-Collas, M., 2016. Exploring methods for predicting multiple pressures on  
1619 ecosystem recovery: A case study on marine eutrophication and fisheries. *Continental*  
1620 *and Shelf Research* 121, 48-60.
- 1621 Vahteri, P., Vuorinen, I., 2016. Continued decline of the bladderwrack, *Fucus*  
1622 *vesiculosus*, in the Archipelago Sea, northern Baltic proper. *Boreal Environment*  
1623 *Research* 21, 373-386.
- 1624 Vahteri, P., Mäkinen, A., Salovius, S., Vuorinen, I., 2000. Are drifting algal mats  
1625 conquering the bottom of the Archipelago Sea, SW Finland? *Ambio* 29, 338-343.
- 1626 Vanhatalo, J., Veneranta, L., Hudd, R., 2012. Species distribution modeling with  
1627 Gaussian processes: A case study with the youngest stages of sea spawning whitefish  
1628 (*Coregonus lavaretus* L. s.l.) larvae. *Ecological Modelling* 228, 49-58.
- 1629 Vasconcelos, R.P., Eggleston, D.B., Le Pape, O., Tulp, I., 2014. Patterns and processes of  
1630 habitat-specific demographic variability in exploited marine species. *ICES Journal of*  
1631 *Marine Science* 71, 638-647.

- 1632 Vetemaa, M., Eschbaum, R., Verliin, A., Albert, A., Eero, M., Lillemägi, R., Pihlak, M.,  
1633 Saat, T., 2006. Annual and seasonal dynamics of fish in the brackish-water Matsalu Bay,  
1634 Estonia.  
1635 Ecology of Freshwater Fish 15, 211-220.
- 1636 Vogt, H., Schramm, W., 1991. Conspicuous decline of *Fucus* in Kiel Bay (western  
1637 Baltic): what are the causes? Marine Ecology Progress Series 69, 189-194.
- 1638 Wastie, J., 2014. Assessing the importance of freshwater tributary systems for the  
1639 recruitment of Eurasian perch (*Perca fluviatilis*) in Baltic Sea coastal ecosystems. MSc-  
1640 thesis, Swedish University of Agricultural Sciences, 30 pp.
- 1641 Weinstein, M.P., Balletto, J.H., 1999. Does the common reed, *Phragmites australis*,  
1642 affect essential fish habitat? Estuaries 22, 793-802.
- 1643 Wennhage, H., Pihl, L., 2002. Fish feeding guilds in shallow rocky and soft bottom  
1644 areas on the Swedish west coast. Journal of Fish Biology 61, 207-228.
- 1645 Węśławski, J.M., Warzocha, J., Wiktor, J., Urbański, J., Bradtke, K., Kryla, L., Tatarek, A.,  
1646 Kotwicki, L., Piwowarczyk, J., 2009. Biological valorisation of the southern Baltic Sea  
1647 (Polish Exclusive Economic Zone). Oceanologia 51, 415-435.
- 1648 Węśławski, J.M., Kryla-Straszewska, L., Piwowarczyk, J., Urbański, J., Warzocha, J.,  
1649 Kotwicki, L., Włodarska-Kowalczyk, M., Wiktor, J., 2013. Habitat modelling limitations –  
1650 Puck Bay, Baltic Sea – a case study. Oceanologia 55, 167-183.
- 1651 Westin, L., Limburg, K.E., 2002. Newly discovered reproductive isolation reveals  
1652 sympatric populations of *Esox lucius* in the Baltic. Journal of Fish Biology 61, 1647-1652.
- 1653 Widdows, J., 1991. Physiological ecology of mussel larvae. Aquaculture 94, 147-163.
- 1654 Winkler, H.M., 1996. Peculiarities of fish communities in estuaries of the Baltic Sea;  
1655 the Darss-Zingst-Bodden Chain as an example. Oceanographic Literature Review 12,  
1656 1258.

Yáñez-Arancibia, A., Day, J.W., Reyes, E., 2013. Understanding the coastal ecosystem-based management approach in the Gulf of Mexico. *Journal of Coastal Research* 63, 243-261.

**Web-links:**

[http://ec.europa.eu/environment/nature/conservation/species/redlist/index\\_en.htm](http://ec.europa.eu/environment/nature/conservation/species/redlist/index_en.htm)

(page visited 27/11/2017)

<http://helcom.fi/helcom-at-work/projects/holas-ii/> (page visited 27/11/2017)

<http://helcom.fi/helcom-at-work/projects/tapas> (page visited 27/11/2017)

<http://www.marlin.ac.uk/> (page visited 27/11/2017)

<http://paikkatieto.ymparisto.fi/velmu/> (page visited 27/11/2017)

<http://www.helcom.fi/> (page visited 27/11/2017)

<https://www.nobanis.org/marine-identification-key/fish/fish-start/fish->

[key/neogobius-melanostomus/](https://www.nobanis.org/marine-identification-key/fish/fish-start/fish-key/neogobius-melanostomus/) (page visited 27/11/2017)



| Direct evidence  |                                     |  |  |   |
|--|-------------------------------------|--|--|---|
| Fish species   | Area                                | Studied topic(s)   | Central findings   | Reference(s)  |
| Perch ( <i>Perca fluviatilis</i> ) and pikeperch ( <i>Sander lucioperca</i> )                              | Sweden and Finland                  | Species distribution modelling was used on coastal data from twelve archipelago areas where the distribution of nursery habitats for perch and pikeperch was related to the size of adult populations. | Habitat availability explains almost half of the variation in population size and indicates a crucial role in limiting adult stock sizes.  | Sundblad et al. 2014  |
| Whitefish ( <i>Coregonus lavaretus</i> )   | Gulf of Bothnia, Sweden and Finland | Species distribution modelling was used on coastal data on whitefish to evaluate relationships between variables describing EFH and larval production.   | Metrics describing EFH and their current level of human impact are the most important ones for the abundance of whitefish larvae.  | Vanhatalo et al. 2012   |
| Perch, pikeperch, Baltic herring ( <i>Clupea harengus membras</i> ) and sprat ( <i>Osmerus eperlanus</i> ) | Finnish coast                       | Species distribution modelling was used on larval survey data of a number of fish species to assess the most important reproduction habitats.  | Identification of highly effective spawning areas, i.e. that production of fish stocks can be concentrated to very limited areas compared to the total suitable production areas that are available.               | Kallasvuo et al. 2017   |
| Cod ( <i>Gadus morhua</i> )*   | Baltic Proper                       | Various statistical models were used for the determination of relationships between the volume of EFH (coastal and non-coastal) available for Baltic cod and processes affecting adult stock size.     | Positive relationships exist between the volume of EFH and cod reproduction (and thus the adult stock size) as well as between habitat availability for juvenile cod (nursery areas) and density-dependent growth. | MacKenzie et al. 2000, Cardinale and Arrhenius 2000, Hinrichsen et al. 2017 |
| Flounder ( <i>Platichthys flesus</i> )*  | Baltic Proper                       | Spawning area availability of pelagic spawning flounder through time was quantified by species distribution modelling, and related to larval production and adult stock sizes.                         | Decreases in spawning habitat availability have been accompanied by a decrease in larval production as well as a decrease in adult stock sizes.  | Ustups et al. 2013, Orio et al. 2017  |
| Indirect evidence  |                                     |  |  |   |
| Fish species   | Area                                | Studied topic(s)   | Central findings   | Reference(s)  |
| Perch  | Sweden and Finland                  | Investigation of how coastal spawning habitats for perch are dependent on the type of substrate.   | Vegetated substrates providing rigidity and structural complexity are preferred by the perch. Also, shallow depths and sheltered areas are preferred characteristics.  | Snickars et al. 2010  |
| Juvenile fish  | Sweden and Finland                  | Relationships between fish reproduction data and the quality (species composition) of macrophyte communities on shallow soft bottoms were investigated   | Investigated bays that are dominated by stress sensitive macrophyte species are important nursery areas for fish.  | Hansen and Snickars 2014  |
| Pike ( <i>Esox lucius</i> )  | Southern Finland                    | Habitat choice and survival of pike in filamentous algae and in bladder-wrack were   | In the presence of predators, pike larvae prefer and also survive better in filamentous algae  | Engström-Öst et al. 2007  |

|  |                          |   |  |   |
|--|--------------------------|---|--|---|
|  |                          | tested experimentally.  | than in bladderwrack.  |   |
| Pike   | Southern Finland         | The performance of larval pike under the influence of turbidity induced by phytoplankton was investigated experimentally.   | Larval weight of pike is lower in turbid water, despite that pike larvae here spend less time in vegetation and attack more prey.  | Engström-Öst and Mattila 2008                     |
| Commercial fish species  | Entire Finnish coastline | Relationships between many environmental variables (N, P, chlorophyll <i>a</i> , duration of ice coverage in winter, shore density in the area and salinity) and the CPUE (of reported commercial catches) were investigated. | Shore density, corresponding closely to the availability of EFH, is an important factor for all species, although the strongest effects occur for pike.  | Uusitalo et al. 2012                              |
| Flounder   | Southern Finland         | Fishery-independent data on adult flounder as well as historical and present-state data on juveniles in shallow coastal areas were utilized to study relationships between EFH and the production of flounder.                | Increased coverage of filamentous algae correlates with a pronounced decrease in the abundance of juvenile flounder. A simultaneous decrease in the abundance of adult flounders indicates that the declined EFH availability for juveniles acts as a bottleneck for the population. | Jokinen et al. 2015, 2016                         |
| Pikeperch  | Germany                  | Investigation of pikeperch spawning in inner coastal waters of salinities around 5-6.   | Coastal EFH of lower salinities are the base for nearly 40% of the total annual catch of pikeperch in waters with higher salinities (around 10).   | Winkler 1996, Winkler et al. unpubl.              |
| Pike   | Southeastern Sweden      | The recruitment of pike was studied in coastal wetlands restored in different ways.   | In restored wetlands with temporally flooded terrestrial vegetation, juvenile pike migration increase from a few thousand individuals in previous years to >100,000 individuals afterwards.  | Nilsson et al. 2014                               |
| Pike   | Swedish east coast       | The relative importance of fresh and brackish water recruitment areas (spawning habitat preferences) for pike was examined through the use of otolith Sr:Ca profiles.   | For pike, 20% hatches in brackish water in the Forsmark area at the 60° N latitude and 80% hatches in brackish water in the Kalmar Sound at the 56° N latitude.  | Engstedt et al. 2010                              |
| Pike, whitefish, burbot ( <i>Lota lota</i> ) and ide ( <i>Leuciscus idus</i> ) | Estonia                  | The relative importance of fresh and brackish water recruitment areas (spawning habitat preferences) was examined for brackish water fish populations through the use of otolith Sr:Ca profiles.                              | The relative importance of coastal wetlands and river-mouths as spawning grounds compared to brackish water areas is demonstrated. There are indications that brackish water spawning is becoming rarer.   | Rohtla et al. 2012, 2014, 2015, 2017, Rohtla 2015 |

**Table 1. Direct and indirect evidence from the Baltic Sea with regard to the effects of EFH on fish population size. \*Note that the results for cod and flounder in the Baltic Proper are not strictly coastal (see text for more details).**